

Spin hydrodynamics

Dirk H. Rischke

with

David Wagner, Nora Weickgenannt, Enrico Speranza


based on

arxiv:2203.04766 [nucl-th], arXiv:2208.01955 [nucl-th]

HFHF Theory Retreat 2022

Castiglione della Pescaia, Sep. 12 – 16, 2022



Λ polarization along angular-momentum direction ("global polarization")


L. Adamczyk et al. (STAR), Nature 548 (2017) 62

→ QGP is "most vortical fluid ever observed"
 $\omega \simeq (9 + 1) \times 10^{21} \text{ s}^{-1}$



For comparison:

- Great Red Spot of Jupiter $\omega \simeq 10^{-4} \text{ s}^{-1}$
- turbulent flow in superfluid He-II $\omega \sim 150 \text{ s}^{-1}$
- superfluid nanodroplets $\omega \sim 10^7 \text{ s}^{-1}$

Assuming local equilibrium on freeze-out hypersurface,
hydrodynamics describes global polarization quite well . . .



$$\Pi^\mu \sim \epsilon^{\mu\nu\rho\sigma} k_\nu \int_{\Sigma_{\text{f.o.}}} d\Sigma \cdot k \varpi_{\rho\sigma} f_{0k}$$

where $\varpi_{\rho\sigma} \equiv -\frac{1}{2} (\partial_\rho \beta_\sigma - \partial_\sigma \beta_\rho)$ thermal vorticity,
 $\beta^\mu \equiv u^\mu / T$, f_{0k} local-equilibrium distribution function

I. Karpenko, F. Becattini, NPA 967 (2017) 764

. . . but fails to describe azimuthal-angle dependence of polarization along the beam direction ("local longitudinal polarization")

F. Becattini, M.A. Lisa, Ann. Rev. Nucl. Part. Sci. 70 (2020) 395



Recently, further (dissipative?) contributions to the polarization were found


F. Becattini, M. Buzzegoli, G. Inghirami, I. Karpenko, A. Palermo, PRL 127 (2021) 272302

$$\Pi^\mu \sim \epsilon^{\mu\nu\rho\sigma} k_\nu \int_{\Sigma_{\text{f.o.}}} d\Sigma \cdot k \left(\varpi_{\rho\sigma} + 2\hat{t}_\rho \xi_{\sigma\lambda} \frac{k^\lambda}{k_0} \right) f_{0k}$$



where $\xi_{\sigma\lambda} \equiv \frac{1}{2} (\partial_\sigma \beta_\lambda + \partial_\lambda \beta_\sigma)$ thermal shear tensor

(see also S.Y.F. Liu, Y. Yin, JHEP 07 (2021) 188)

→ this does not quite do the job ...



... but neglecting temperature gradients on $\Sigma_{\text{f.o.}}$ does!



Observations:

- derivation of formula for polarization ($\Pi^\mu \sim \varpi_{\rho\sigma}$) is strictly valid only for rotating global-equilibrium state
F. Becattini, V. Chandra, L. Del Zanna, E. Grossi, Annals Phys. 338 (2013) 32
⇒ application of formula requires (infinitely) fast equilibration of spin degrees of freedom (relative to timescale of collision)
- dissipative effects influence (almost) all other observables in a heavy-ion collision, even appear explicitly in new term $\sim \xi_{\sigma\lambda}$ in formula for polarization

⇒ Questions:

- (I) How fast do spin degrees of freedom equilibrate?
- (II) How is polarization influenced by dissipative effects?

⇒ requires a theory of second-order dissipative spin hydrodynamics!

N. Weickgenannt, D. Wagner, E. Speranza, DHR,
arxiv:2203.04766 [nucl-th], arXiv:2208.01955 [nucl-th]

Remark: for tensor polarization and spin-1 particles, see D. Wagner's talk

Particle number conservation

$$\partial_\mu N^\mu = 0$$

where N^μ particle four-current

Energy-momentum conservation

$$\partial_\mu T^{\mu\nu} = 0$$

where $T^{\mu\nu}$ energy-momentum tensor

Angular-momentum conservation

$$\partial_\mu J^{\mu,\nu\lambda} = 0$$

where $J^{\mu,\nu\lambda}$ angular-momentum tensor

with $J^{\mu,\nu\lambda} \equiv x^\nu T^{\mu\lambda} - x^\lambda T^{\mu\nu} + \hbar S^{\mu,\nu\lambda}$ and energy-momentum conservation:

Equation of motion for spin tensor

$$\hbar \partial_\mu S^{\mu,\nu\lambda} = T^{[\lambda\nu]}$$

where $a^{[\lambda} b^{\nu]} \equiv a^\lambda b^\nu - a^\nu b^\lambda$

Particle number conservation

$$\partial_\mu N^\mu = 0$$

1 equation, 4 unknowns

Energy-momentum conservation

$$\partial_\mu T^{\mu\nu} = 0$$

4 equations, (at least) 10 unknowns

Angular-momentum conservation

$$\partial_\mu J^{\mu,\nu\lambda} = 0$$

with $J^{\mu,\nu\lambda} \equiv x^\nu T^{\mu\lambda} - x^\lambda T^{\mu\nu} + \hbar S^{\mu,\nu\lambda}$ and energy-momentum conservation:

Equation of motion for spin tensor

$$\hbar \partial_\mu S^{\mu,\nu\lambda} = T^{[\lambda\nu]}$$

6 equations, 24 unknowns

where $a^{[\lambda} b^{\nu]} \equiv a^\lambda b^\nu - a^\nu b^\lambda$

Assume local equilibrium

Single-particle distribution function (to order $\mathcal{O}(\hbar)$)



$$f_{\text{eq}, \mathbf{k}} = f_0 \mathbf{k} \left(1 + \frac{\hbar}{4} \Omega_{\mu\nu} \Sigma_{\mathbf{k}}^{\mu\nu} \right), \quad f_0 = \exp(\alpha - \beta \cdot k)$$

where

- $\alpha \equiv \frac{\mu}{T}$ Lagrange multiplier for particle-number conservation
- $\beta^\mu \equiv \frac{u^\mu}{T}$ Lagrange multiplier for energy-momentum conservation, u^μ fluid 4-velocity
- $\Omega_{\mu\nu}$ spin potential, Lagrange multiplier for angular-momentum conservation
- $\Sigma_{\mathbf{k}}^{\mu\nu} \equiv -\frac{1}{m} \epsilon^{\mu\nu\alpha\beta} k_\alpha \mathbf{s}_\beta$ dipole-moment tensor of particle with mass m , (on-shell) 4-momentum k_α , and spin 4-vector \mathbf{s}_β

Assume local equilibrium

Single-particle distribution function (to order $\mathcal{O}(\hbar)$)



$$f_{\text{eq}, \mathbf{k}} = f_0 \mathbf{k} \left(1 + \frac{\hbar}{4} \Omega_{\mu\nu} \Sigma_{\mathbf{k}}^{\mu\nu} \right), \quad f_0 \mathbf{k} = \exp(\alpha - \beta \cdot \mathbf{k})$$

where

- $\alpha \equiv \frac{\mu}{T}$ Lagrange multiplier for particle-number conservation
1 parameter
- $\beta^\mu \equiv \frac{u^\mu}{T}$ Lagrange multiplier for energy-momentum conservation, u^μ fluid 4-velocity
4 parameters
- $\Omega_{\mu\nu}$ spin potential, Lagrange multiplier for angular-momentum conservation
6 parameters
- $\Sigma_{\mathbf{k}}^{\mu\nu} \equiv -\frac{1}{m} \epsilon^{\mu\nu\alpha\beta} k_\alpha \mathbf{s}_\beta$ dipole-moment tensor of particle with mass m ,
(on-shell) 4-momentum k_α , and spin 4-vector \mathbf{s}_β

Extend phase space by spin degrees of freedom (for more details, see D. Wagner's talk):

$$dK \equiv \frac{d^3 k}{(2\pi)^3} \longrightarrow d\Gamma \equiv dK \, dS(k)$$

with $dS(k) \equiv \sqrt{\frac{k^2}{3\pi^2}} d^4 s \delta(k \cdot s) \delta(s \cdot s + 3),$

such that $\int dS(k) = 2, \int dS(k) s^\mu = 0, \int dS(k) s^\mu s^\nu = -2 \left(g^{\mu\nu} - \frac{k^\mu k^\nu}{k^2} \right)$

Fluid-dynamical currents

$$\begin{aligned} N^\mu &\equiv \langle k^\mu \rangle \\ T^{\mu\nu} &\equiv \langle k^\mu k^\nu \rangle + \mathcal{O}(\hbar^2) \\ S^{\mu,\nu\lambda} &\equiv \left\langle k^\mu \left(\frac{1}{2} \Sigma_{\mathbf{k}s}^{\nu\lambda} - \frac{\hbar}{4m^2} k^{[\nu} \partial^{\lambda]} \right) \right\rangle + \mathcal{O}(\hbar^2) \end{aligned}$$

where $\langle \cdots \rangle \equiv \int d\Gamma \cdots f(x, k, s)$

\Rightarrow for $\langle \cdots \rangle_{\text{eq}} \equiv \int d\Gamma \cdots f_{\text{eq}, \mathbf{k}s} \Rightarrow$ equations of motion are closed!

see, e.g., W. Florkowski, A. Kumar, R. Ryblewski, Prog. Part. Nucl. Phys. 108 (2019) 103709

Dissipative spin hydrodynamics

- ⇒ provide additional equations of motion
- ⇒ start from underlying microscopic theory, apply method of moments
see, e.g., G.S. Denicol, H. Niemi, E. Molnár, DHR, PRD 85 (2012) 114047

From equation of motion for the Wigner function, derive to first order in \hbar :

N. Weickgenannt, E. Speranza, X.-I. Sheng, Q. Wang, DHR,

PRL 127 (2021) 052301, PRD 104 (2021) 016022 for more details, see D. Wagner's talk

Boltzmann equation for spin-1/2 particles with nonlocal collision term

$$\begin{aligned} k \cdot \partial f(x, k, \mathfrak{s}) &= \mathfrak{C}[f] \\ \mathfrak{C}[f] &= \int d\Gamma_1 d\Gamma_2 d\Gamma' \mathcal{W}_{kk' \rightarrow k_1 k_2}^{\mathfrak{s}\mathfrak{s}' \rightarrow \mathfrak{s}_1 \mathfrak{s}_2} [f(x + \Delta_1, k_1, \mathfrak{s}_1) f(x + \Delta_2, k_2, \mathfrak{s}_2) \\ &\quad - f(x + \Delta, k, \mathfrak{s}) f(x + \Delta', k', \mathfrak{s}')] \end{aligned}$$

where nonlocal position shift $\Delta^\mu = -\frac{\hbar}{2m(k \cdot \hat{t} + m)} \epsilon^{\mu\nu\alpha\beta} k_\nu \hat{t}_\alpha \mathfrak{s}_\beta$

- ↔ Berry connection! see, e.g., M. Stone, V. Dwivedi, T. Zhou, PRD 91 (2015) 025004

Note: $\Delta \sim \frac{\hbar}{m}$ Compton wavelength!

Nonlocal collisions: allow mutual conversion of orbital angular momentum and spin

Usually, local-equilibrium distribution function f_{eq} is defined by condition

Local equilibrium

$$\mathfrak{C}[f_{\text{eq}}] \equiv 0$$

However, as shown in N. Weickgenannt, E. Speranza, X.-I. Sheng, Q. Wang, DHR, PRL 127 (2021) 052301, PRD 104 (2021) 016022, nonlocal collision term vanishes only in

Global equilibrium

$$\begin{aligned}\alpha &= \text{const.} \\ \partial^\mu \beta^\nu + \partial^\nu \beta^\mu &= 0 \\ \Omega_{\mu\nu} &= \varpi_{\mu\nu} = \text{const.}\end{aligned}$$

- ⇒ appears too restrictive: nonlocality $\Delta \lesssim r_{\text{int}} \ll \lambda_{\text{mfp}} \ll L_{\text{hydro}}$
- ⇒ nonlocality scale Δ much smaller than hydrodynamic scale L_{hydro}

Generalized local equilibrium



$$\mathfrak{C}[f_{\text{eq}}] \sim \mathcal{O}(\Delta/L_{\text{hydro}})$$

Define hydrodynamic scale L_{hydro} by

$$\frac{1}{m} k \cdot \partial f_{\text{eq}, \mathbf{k}} \sim \frac{1}{L_{\text{hydro}}} f_{\text{eq}, \mathbf{k}}$$

$$\begin{aligned}\partial_\mu \alpha &\sim \mathcal{O}(L_{\text{hydro}}^{-1}) \\ \partial^\mu \beta^\nu + \partial^\nu \beta^\mu &\sim \mathcal{O}((k L_{\text{hydro}})^{-1}) \\ \partial_\lambda \Omega_{\mu\nu} &\sim \mathcal{O}(L_{\text{hydro}}^{-1})\end{aligned}$$

using conservation of total angular momentum $J^{\mu\nu} \equiv \Delta^{[\mu} k^{\nu]} + \frac{\hbar}{2} \Sigma_{\mathbf{k}}^{\mu\nu}$ in binary collisions, order $\mathcal{O}(\hbar)$ contribution to nonlocal collision term:

$$\sim \frac{\hbar}{4} \Omega_{\mu\nu} \Sigma_{\mathbf{k}}^{\mu\nu} + \Delta^{\mu} \partial_\mu (\alpha - \beta^\nu k^\nu) = \frac{1}{2} \Delta^{[\mu} k^{\nu]} (\varpi_{\mu\nu} - \Omega_{\mu\nu}) + \mathcal{O}(\Delta/L_{\text{hydro}})$$

for generalized local equilibrium: $\Omega_{\mu\nu} \equiv \varpi_{\mu\nu} + \mathcal{O}((k L_{\text{hydro}})^{-1})$

consistent, as in global equilibrium $L_{\text{hydro}} \rightarrow \infty$ and thus $\Omega_{\mu\nu} \rightarrow \varpi_{\mu\nu}$

Usually, all gradients of fluid-dynamical quantities are $\mathcal{O}(L_{\text{hydro}}^{-1})$

However, global-equilibrium conditions do not restrict value of thermal vorticity $\varpi_{\mu\nu}$
see, e.g., F. Becattini, L. Tinti, Annals Phys. 325 (2010) 1566

⇒ vorticity does not follow usual power counting, $\varpi_{\mu\nu} \not\sim \mathcal{O}((kL_{\text{hydro}})^{-1})$

⇒ define scale ℓ_{vort} set by vorticity: $\varpi_{\mu\nu} \sim \mathcal{O}((k\ell_{\text{vort}})^{-1})$

In principle, ℓ_{vort} can take any value from $\ell_{\text{vort}} \ll L_{\text{hydro}}$ to $\ell_{\text{vort}} \sim L_{\text{hydro}}$

However, in order for \hbar -expansion to apply: $\hbar\Omega_{\mu\nu}\Sigma_{\mathbf{k}s}^{\mu\nu} \sim \frac{\hbar}{m}\varpi_{\mu\nu}\epsilon^{\mu\nu\alpha\beta}k_{\alpha}s_{\beta} \sim \frac{\Delta}{\ell_{\text{vort}}} \ll 1$

⇒ ℓ_{vort} cannot be arbitrarily small (as in global equilibrium)

⇒ remember $\Delta \lesssim r_{\text{int}} \ll \lambda_{\text{mfp}} \ll L_{\text{hydro}}$ ⇒ ℓ_{vort} could be as small as $\lambda_{\text{mfp}}!$

⇒ for the sake of simplicity assume

$$\frac{\Delta}{\ell_{\text{vort}}} \sim \frac{\lambda_{\text{mfp}}}{L_{\text{hydro}}} \equiv Kn \ll 1$$

Extend method of moments developed in G.S. Denicol, H. Niemi, E. Molnár, DHR, PRD 85 (2012) 114047 by spin degrees of freedom

Single-particle distribution function

$$f(x, k, \mathbf{s}) = f_{\text{eq}, \mathbf{k}\mathbf{s}} + \delta f_{\mathbf{k}\mathbf{s}}$$

$$\delta f_{\mathbf{k}\mathbf{s}} = f_{0\mathbf{k}} \sum_{\ell=0}^{\infty} \sum_{n \in \mathbb{S}_{\ell}} \mathcal{H}_{\mathbf{k}n}^{(\ell)} \left[\rho_n^{\mu_1 \dots \mu_{\ell}} - \tau_n^{\langle \mu \rangle, \mu_1 \dots \mu_{\ell}} \left(g_{\mu\nu} - \frac{k_{\langle \mu \rangle} u_{\nu}}{E_{\mathbf{k}}} \right) \mathbf{s}^{\nu} \right] k_{\langle \mu_1} \dots k_{\mu_{\ell} \rangle}$$

where

- $k_{\langle \mu_1 \dots k_{\mu_{\ell}} \rangle}$ irreducible tensors
- $\rho_n^{\mu_1 \dots \mu_{\ell}} \equiv \langle E_{\mathbf{k}}^n k^{\langle \mu_1} \dots k^{\mu_{\ell}} \rangle_{\delta}$ irreducible moments, $\langle \dots \rangle_{\delta} \equiv \langle \dots \rangle - \langle \dots \rangle_{\text{eq}}$
- $\tau_n^{\mu, \mu_1 \dots \mu_{\ell}} \equiv \langle \mathbf{s}^{\mu} E_{\mathbf{k}}^n k^{\langle \mu_1} \dots k^{\mu_{\ell}} \rangle_{\delta}$ spin moments
- $E_{\mathbf{k}} \equiv k \cdot u$ energy of particle in fluid rest frame
- $A^{\langle \mu \rangle} \equiv \Delta^{\mu\nu} A_{\nu}$, with $\Delta^{\mu\nu} \equiv g^{\mu\nu} - u^{\mu} u^{\nu}$ projector onto 3-space orthogonal to u^{μ}
- $A^{\langle \mu_1 \dots \mu_{\ell} \rangle} \equiv \Delta_{\nu_1 \dots \nu_{\ell}}^{\mu_1 \dots \mu_{\ell}} A^{\nu_1 \dots \nu_{\ell}}$,
 $\Delta_{\nu_1 \dots \nu_{\ell}}^{\mu_1 \dots \mu_{\ell}}$ symmetric, traceless rank- 2ℓ projection operators built from $\Delta^{\mu\nu}$
- $\mathcal{H}_{\mathbf{k}n}^{(\ell)}$ polynomials in $E_{\mathbf{k}}$ of rank N_{ℓ}

Inserting $f(x, k, \mathfrak{s}) = f_{\text{eq}, \mathbf{k}\mathfrak{s}} + \delta f_{\mathbf{k}\mathfrak{s}}$ into definition of spin tensor

Spin tensor



$$S^{\mu, \nu, \lambda} = u^\mu \tilde{\mathfrak{N}}^{\nu, \lambda} + \tilde{\mathfrak{P}}^{\langle \mu \rangle, \nu, \lambda} + u_\alpha \tilde{\mathfrak{H}}^{\mu, \nu, \lambda, \alpha} + u^\mu \tilde{\mathfrak{H}}_\alpha^{\nu, \lambda, \alpha} + \tilde{\mathfrak{Q}}^{\mu, \nu, \lambda} - \frac{\hbar}{4m^2} \partial^{[\lambda} T^{\nu]\mu}$$

where

- $\tilde{\mathfrak{N}}^{\nu, \lambda} \equiv \epsilon^{\nu, \lambda, \alpha, \beta} \mathfrak{N}_{\alpha\beta}$,
with $\mathfrak{N}^{\alpha\beta} \equiv -\frac{1}{2m} u^\alpha \left[\left\langle E_{\mathbf{k}}^2 \mathfrak{s}^\beta \right\rangle_{\text{eq}} + \tau_2^\beta \right]$ spin energy tensor
- $\tilde{\mathfrak{P}}^{\mu, \nu, \lambda} \equiv \epsilon^{\mu, \nu, \lambda, \alpha} \mathfrak{P}_\alpha$,
with $\mathfrak{P}^\alpha \equiv -\frac{1}{6m} \left[\left\langle (m^2 - E_{\mathbf{k}}^2) \mathfrak{s}^\alpha \right\rangle_{\text{eq}} + m^2 \tau_0^\alpha - \tau_2^\alpha \right]$ spin pressure vector
- $\tilde{\mathfrak{H}}^{\mu, \nu, \lambda, \alpha} \equiv \epsilon^{\nu, \lambda, \alpha, \beta} \mathfrak{H}^\mu{}_\beta$,
with $\mathfrak{H}^{\mu\beta} \equiv -\frac{1}{2m} \left[\left\langle E_{\mathbf{k}} k^{\langle \mu} \mathfrak{s}^{\beta \rangle} \right\rangle_{\text{eq}} + \tau_1^{\beta, \mu} \right]$ spin diffusion tensor
- $\tilde{\mathfrak{Q}}^{\mu, \nu, \lambda} \equiv \epsilon^{\nu, \lambda, \alpha, \beta} \mathfrak{Q}^\mu{}_{\alpha\beta}$,
with $\mathfrak{Q}^{\mu\alpha\beta} \equiv -\frac{1}{2m} \left[\left\langle k^{\langle \mu} k^{\alpha \rangle} \mathfrak{s}^\beta \right\rangle_{\text{eq}} + \tau_0^{\beta, \mu\alpha} \right]$ spin stress tensor

Define local-equilibrium state via

Landau matching conditions

$$\begin{aligned} N^\mu u_\mu &= N_{\text{eq}}^\mu u_\mu \\ T^{\mu\nu} u_\nu &= T_{\text{eq}}^{\mu\nu} u_\nu \\ J^{\mu,\nu\lambda} u_\mu &= J_{\text{eq}}^{\mu,\nu\lambda} u_\mu \end{aligned}$$

- ⇒ relates α , β^μ , and $\Omega_{\mu\nu}$ in $f_{\text{eq},\mathbf{k}}$ to fluid-dynamical variables
- ⇒ determine α , β^μ , and $\Omega_{\mu\nu}$ via conservation laws!
- ⇒ still need to derive equations of motion for dissipative currents Π , n^μ , and $\pi^{\mu\nu}$, as well as spin moments τ_0^α , τ_2^α , $\tau_1^{\beta,\mu}$, and $\tau_0^{\beta,\mu\alpha}$ appearing in $S^{\mu,\lambda\nu}$

Equations of motion for standard dissipative currents Π , n^μ , and $\pi^{\mu\nu}$

- ⇒ see G.S. Denicol, H. Niemi, E. Molnár, DHR, PRD 85 (2012) 114047
- ⇒ relaxation-type equations, e.g., $\dot{\Pi} + \frac{1}{\tau_\Pi} \Pi = \dots$, with $\dot{\Pi} \equiv u^\mu \partial_\mu \Pi$

Take spin moments of Boltzmann equation:

Equations of motion for spin moments

$$\Rightarrow \dot{\tau}_n^{\langle\mu\rangle,\langle\mu_1\cdots\mu_\ell\rangle} - \mathfrak{C}_{n-1}^{\langle\mu\rangle,\mu_1\cdots\mu_\ell} = \dots$$

$$\mathfrak{C}_{n-1}^{\mu,\mu_1\cdots\mu_\ell} = \int d\Gamma E_{\mathbf{k}}^{n-1} k^{\langle\mu_1} \cdots k^{\mu_\ell\rangle} \mathfrak{s}^\mu \mathfrak{C}[f]$$

Linearized collision integral

$$\mathfrak{C}_{n-1}^{\mu,\mu_1\cdots\mu_\ell} = \mathfrak{C}_{n-1,\text{local}}^{\mu,\mu_1\cdots\mu_\ell} + \mathfrak{C}_{n-1,\text{nonlocal}}^{\mu,\mu_1\cdots\mu_\ell}$$

$$\mathfrak{C}_{n-1,\text{local}}^{\mu,\mu_1\cdots\mu_\ell} = - \sum_{r \in \mathbb{S}_\ell} B_{nr}^{(\ell)} \tau_r^{\mu,\mu_1\cdots\mu_\ell}$$

$$\mathfrak{C}_{n-1,\text{nonlocal}}^{\mu,\mu_1\cdots\mu_\ell} = \int d\Gamma_1 d\Gamma_2 d\Gamma d\Gamma' \mathcal{W}_{\mathbf{k}\mathbf{k}' \rightarrow \mathbf{k}_1 \mathbf{k}_2}^{\mathfrak{s}\mathfrak{s}' \rightarrow \mathfrak{s}_1 \mathfrak{s}_2} E_{\mathbf{k}}^{n-1} f_{0\mathbf{k}} f_{0\mathbf{k}'}$$

$$\times k^{\langle\mu_1} \cdots k^{\mu_\ell\rangle} \mathfrak{s}^\mu \left[\frac{\hbar}{4} (\varpi_{\alpha\beta} - \Omega_{\alpha\beta}) \Sigma_{\mathbf{k}\mathfrak{s}}^{\alpha\beta} + \xi_{\alpha\beta} \Delta^\alpha k^\beta \right]$$

- $\mathfrak{C}_{n-1,\text{local}}^{\mu,\mu_1\cdots\mu_\ell} \Rightarrow$ inverting $B_{nr}^{(\ell)}$ yields relaxation times
- $\mathfrak{C}_{n-1,\text{nonlocal}}^{\mu,\mu_1\cdots\mu_\ell} \Rightarrow$ gives rise to Navier-Stokes terms

Infinite set of moment equations needs to be truncated

⇒ lowest-order truncation:

14 standard fluid-dynamical moments + 24 moments for components of spin tensor

⇒ (14+24)-moment approximation

Independent spin moments: $p^\mu \equiv \tau_0^{\langle\mu\rangle}$, $j^{\mu\nu} \equiv \tau_1^{\langle\mu\rangle,\nu} + \tau_1^{\langle\nu\rangle,\mu}$, $q^{\mu\nu\lambda} \equiv \tau_0^{\langle\mu\rangle,\nu\lambda}$


3 + 6 + 15 components

Equations of motion for independent spin moments

$$\Rightarrow \begin{aligned} \tau_p \dot{p}^{\langle\mu\rangle} + p^\mu &\sim \epsilon^{\mu\nu\alpha\beta} (\varpi_{\alpha\beta} - \Omega_{\alpha\beta}) u_\nu + \dots \\ \tau_j \dot{j}^{\langle\mu\rangle\langle\nu\rangle} + j^{\mu\nu} &\sim \dots \\ \tau_q \dot{q}^{\langle\mu\rangle\langle\nu\lambda\rangle} + q^{\mu\nu\lambda} &\sim \xi_\alpha^{\langle\nu} \epsilon^{\lambda\rangle\mu\alpha\beta} u_\beta + \dots \end{aligned}$$

for more details, see D. Wagner's talk

Spin relaxation times



- ⇒ spin relaxation times of the same order (but somewhat smaller) than relaxation times for $\Pi, n^\mu, \pi^{\mu\nu}$
- ⇒ spin degrees of freedom equilibrate (i.e., approach their Navier-Stokes values) as fast (or even faster) than $\Pi, n^\mu, \pi^{\mu\nu}$
- ⇒ answers Question (I)

Pauli-Lubanski vector (spin polarization vector!) in Navier-Stokes limit

$$\begin{aligned} \Pi_{\text{NS}}^\mu &\sim \int_{\Sigma_{\text{f.o.}}} d\Sigma \cdot k f_{0k} \left\{ \epsilon^{\mu\nu\rho\sigma} k_\nu \Omega_{\rho\sigma} + \left(\delta_\nu^\mu - \frac{u^\mu k_{\langle\nu\rangle}}{E_k} \right) \right. \\ &\times \left. \left[\kappa_p \epsilon^{\nu\rho\alpha\beta} (\Omega_{\alpha\beta} - \varpi_{\alpha\beta}) u_\rho + \kappa_q \xi_\alpha^{(\rho} \epsilon^{\sigma)\nu\alpha\beta} u_\beta k_{\langle\rho} k_{\sigma\rangle} \right] \right\} \end{aligned}$$

- ⇒ novel dissipative corrections $\sim \Omega_{\alpha\beta} - \varpi_{\alpha\beta}$ and $\xi_{\alpha\beta}$ ⇒ answers Question (II)

- Starting from Boltzmann equation with nonlocal collision term, and using method of moments, derived equations of motion of relativistic second-order dissipative spin hydrodynamics in (14+24)-moment approximation
- Spin degrees of freedom relax as fast as usual dissipative quantities
- Polarization vector is influenced by dissipative corrections
- Need to quantify influence of dissipative corrections on polarization observables!
- Causality and stability of equations of motion of spin hydrodynamics?