

Radiative energy loss and damping effects

Joerg Aichelin, and Marcus Bluhm, and Pol Gossiaux,
and Thierry Gousset (SubaTech)

Sept 7, 2011

Energy loss

Introduction

Energy loss

QCD case

(Q)ED case

Summary

- quark and gluon energy loss is a central topic in ultra-relativistic heavy ion collisions

- collisional and radiative

- Bethe-Heitler regime, LPM effect...

...and damping

radiation and energy loss

Introduction

QCD case

ΔE

in electrodynamics

coherence

LPM effect

radiation spectrum

formation time

competing effects

without damping

with damping


discussion

(Q)ED case

Summary

radiative energy loss of quarks or gluons

an energetic parton in a hot medium ($E \gg T$)



in electrodynamics

Introduction

QCD case

ΔE

in electrodynamics

coherence

LPM effect

radiation spectrum

formation time

competing effects


without damping

with damping

discussion

(Q)ED case

Summary



relativistic electron

■ Bethe-Heitler spectrum

→ radiation loss in matter

$$X_0 = \text{radiation length}$$

length scale on which e^- loses its energy (on average)

coherence

Introduction

QCD case

ΔE

in electrodynamics

coherence

LPM effect

radiation spectrum

formation time

competing effects

without damping


with damping

discussion

(Q)ED case

Summary

to be accommodated with another essential of bremsstrahlung



formation time $<$ radiation length

$$(v \approx c = 1)$$

breaks down at small ω and large E

including multiple scattering: LPM effect

Introduction

QCD case

ΔE


in electrodynamics
coherence

LPM effect

radiation spectrum
formation time
competing effects
without damping
with damping
discussion

(Q)ED case

Summary



e^- multiple scattering speeds up decoherence

\Rightarrow emission process cannot occur at full rate

radiation spectrum

Introduction

QCD case

ΔE

in electrodynamics

coherence

LPM effect

radiation spectrum

formation time

competing effects

without damping

with damping

discussion

(Q)ED case

Summary


$dW/d\omega$ for an energetic parton using the scaling behavior

$$dW/d\omega \propto t_{\text{form}}$$

specifically

$$\frac{dW}{d\omega} = \frac{t_{\text{form}}}{t_{\text{BH}}} \times \frac{dW_{\text{BH}}}{d\omega}$$

$t_{\text{form}}?$ \rightarrow diffraction theory



formation time

Introduction

QCD case

ΔE

in electrodynamics

coherence

LPM effect

radiation spectrum

formation time

competing effects

without damping

with damping

discussion

(Q)ED case

Summary

$$\Delta\varphi(t, R) = \omega(t - t_{\text{form}}) - \vec{k} \cdot \overrightarrow{FR} - \omega(t - 0) + \vec{k} \cdot \overrightarrow{OR}$$

$$\Delta\varphi = -\omega t_{\text{form}} + \underbrace{\vec{k} \cdot \overrightarrow{OF}}_{k_{||} v_{||} t_{\text{form}}}$$

$$k_{||} = n \frac{\omega}{c} \cos \theta, \quad v_{||} = v \cos \theta_s$$

$$|\Delta\varphi| \equiv 1 \rightarrow t_{\text{form}}$$

$$t_{\text{form}} = \frac{1}{\omega |1 - nv \cos \theta_s \cos \theta|}$$

large formation time for ultrarelativistic particles, and small angles, and n close to 1

competing effects

Introduction

QCD case

ΔE

in electrodynamics

coherence

LPM effect

radiation spectrum

formation time

competing effects

without damping

with damping

discussion

(Q)ED case

Summary

1. wave propagation in medium (no damping)

$$n(\omega) = \sqrt{1 - \frac{m_g^2}{\omega^2}}$$

competing effects

Introduction

QCD case

ΔE

in electrodynamics

coherence

LPM effect

radiation spectrum

formation time

competing effects

without damping

with damping

discussion

(Q)ED case

Summary

1. wave propagation in medium (no damping) : n -driven regime ($v \cos \theta_s \rightarrow 1, \theta \rightarrow 0$)

$$t_1 = \frac{1}{\omega(n-1)} \sim \frac{2\omega}{m_g^2}$$

competing effects

Introduction

QCD case

ΔE

in electrodynamics

coherence

LPM effect

radiation spectrum

formation time

competing effects

without damping

with damping

discussion

(Q)ED case

Summary

1. wave propagation in medium (no damping) : n -driven regime ($v \cos \theta_s \rightarrow 1, \theta \rightarrow 0$)

$$t_1 = \frac{1}{\omega(n-1)} \sim \frac{2\omega}{m_g^2}$$

2. (gluon) multiple scattering

$$k_{||}(t) = n\omega(1 - \hat{q}t/\omega^2)$$

competing effects

Introduction

QCD case

ΔE

in electrodynamics

coherence

LPM effect

radiation spectrum

formation time

competing effects

without damping

with damping

discussion

(Q)ED case

Summary

1. wave propagation in medium (no damping) : n -driven regime ($v \cos \theta_s \rightarrow 1, \theta \rightarrow 0$)

$$t_1 = \frac{1}{\omega(n-1)} \sim \frac{2\omega}{m_g^2}$$

2. (gluon) multiple scattering : multiple-scattering-driven regime ($n \rightarrow 1, v \cos \theta_s \rightarrow 1$)

$$t_2 = \frac{1}{\omega(\hat{q} t_2 / \omega^2)} = \sqrt{\frac{\omega}{\hat{q}}}$$

competing effects

Introduction

QCD case

ΔE

in electrodynamics

coherence

LPM effect

radiation spectrum

formation time

competing effects

without damping

with damping

discussion

(Q)ED case

Summary

1. wave propagation in medium (no damping) : n -driven regime ($v \cos \theta_s \rightarrow 1, \theta \rightarrow 0$)

$$t_1 = \frac{1}{\omega(n-1)} \sim \frac{2\omega}{m_g^2}$$

2. (gluon) multiple scattering : multiple-scattering-driven regime ($n \rightarrow 1, v \cos \theta_s \rightarrow 1$)

$$t_2 = \frac{1}{\omega(\hat{q} t_2 / \omega^2)} = \sqrt{\frac{\omega}{\hat{q}}}$$

3. damping

1 and 2

Introduction

QCD case

ΔE

in electrodynamics

coherence

LPM effect

radiation spectrum

formation time

competing effects

without damping


with damping

discussion

(Q)ED case


Summary

$[\log] t_{\text{form}}$



$[\log] \omega$

$[\log] \frac{dW}{d\omega}$



$[\log] \omega$

1 and 2

Introduction

QCD case

ΔE

in electrodynamics

coherence

LPM effect

radiation spectrum

formation time

competing effects


without damping

with damping

discussion

(Q)ED case

Summary



$$t_1 = t_2 \quad \Rightarrow \quad \omega_{\text{LPM}} = \frac{m_g^4}{\hat{q}} \quad \text{at which} \quad t_1 = m_g^2 / \hat{q} = \lambda$$

+ damping

$$\sim e^{-\Gamma t} \Rightarrow \text{damping regime when } t_{\text{form}} \gg 1/\Gamma$$

Introduction

QCD case

ΔE

in electrodynamics

coherence

LPM effect

radiation spectrum

formation time

competing effects

without damping

with damping

discussion

(Q)ED case

Summary

+ damping

Introduction

QCD case

ΔE

in electrodynamics

coherence

LPM effect

radiation spectrum

formation time

competing effects

without damping


with damping

discussion

(Q)ED case

Summary

$\sim e^{-\Gamma t} \Rightarrow$ damping regime when $t_{\text{form}} \gg 1/\Gamma$



+ damping

Introduction

QCD case

ΔE

in electrodynamics

coherence

LPM effect

radiation spectrum

formation time

competing effects

without damping


with damping

discussion

(Q)ED case

Summary

$\sim e^{-\Gamma t} \Rightarrow$ damping regime when $t_{\text{form}} \gg 1/\Gamma$



$$t_2 = 1/\Gamma \Rightarrow \omega_{\text{damp}} = \frac{\hat{q}}{\Gamma^2}$$

discussion

Introduction

QCD case

ΔE

in electrodynamics

coherence

LPM effect

radiation spectrum

formation time

competing effects

without damping

with damping

discussion

(Q)ED case

Summary

■ $\lambda = O(1/(g^2 T))$ and $\Gamma = O(g^2 T)$

$$\frac{1}{\Gamma} \sim \lambda$$

■ is data (jet quenching) compatible with $\Gamma \sim \frac{1}{\lambda}$?

■ quantitatively?

■ microscopic origin of Γ ?

goals

Introduction

QCD case

(Q)ED case

goals

arXiv:1106.2856v1

without damping

with damping

Summary

- confronting the above reasoning with a true calculation
- trace back formation time and damping factor
- compare computed spectrum with the formation-time scaling law

$$\frac{dW}{d\omega} = \frac{t_{\text{form}}}{t_{\text{BH}}} \times \frac{dW_{\text{BH}}}{d\omega}$$

(with ED-type formation times)

from arXiv:1106.2856v1

Introduction

QCD case

(Q)ED case

goals

arXiv:1106.2856v1

without damping

with damping

Summary

energy loss in an absorptive dielectric medium

$$n^2(\omega) = 1 - \frac{m^2}{\omega^2} + 2i\frac{\Gamma}{\omega}$$

using linear response theory \rightarrow mechanical work on charge

$$W = 2 \operatorname{Re} \left(\int d^3\vec{r}' \int d\omega \vec{E}(\vec{r}', \omega) \cdot \vec{j}(\vec{r}', \omega)^* \right)$$

$$\begin{aligned} \frac{d^2W}{dzd\omega} &\simeq -\operatorname{Re} \left(\frac{2i\alpha}{3\pi} \frac{\hat{q}}{E^2} \int_0^\infty d\bar{t} \frac{\omega n^2}{\epsilon} \exp \left[-\omega |n_i| \beta \bar{t} \left(1 - \frac{\hat{q}}{6E^2} \bar{t} \right) \right] \right. \\ &\quad \left. \times \cos(\omega \bar{t}) \exp \left[i\omega n_r \beta \bar{t} \left(1 - \frac{\hat{q}}{6E^2} \bar{t} \right) \right] \right) \end{aligned}$$

from arXiv:1106.2856v1

Introduction

QCD case

(Q)ED case

goals

arXiv:1106.2856v1

without damping

with damping

Summary

energy loss in an absorptive dielectric medium

$$n^2(\omega) = 1 - \frac{m^2}{\omega^2} + 2i\frac{\Gamma}{\omega}$$

using linear response theory \rightarrow mechanical work on charge

$$W = 2 \operatorname{Re} \left(\int d^3\vec{r}' \int d\omega \vec{E}(\vec{r}', \omega) \cdot \vec{j}(\vec{r}', \omega)^* \right)$$

$$\begin{aligned} \frac{d^2W}{dzd\omega} &\simeq -\operatorname{Re} \left(\frac{2i\alpha}{3\pi} \frac{\hat{q}}{E^2} \int_0^\infty d\bar{t} \frac{\omega n^2}{\epsilon} \right. \\ &\quad \left. \times \cos(\omega\bar{t}) \exp \left[i\omega n_r \beta \bar{t} \left(1 - \frac{\hat{q}}{6E^2} \bar{t} \right) \right] \right) \end{aligned}$$

from arXiv:1106.2856v1

Introduction

QCD case

(Q)ED case

goals

arXiv:1106.2856v1

without damping

with damping

Summary

energy loss in an absorptive dielectric medium

$$n^2(\omega) = 1 - \frac{m^2}{\omega^2} + 2i\frac{\Gamma}{\omega}$$

using linear response theory \rightarrow mechanical work on charge

$$W = 2 \operatorname{Re} \left(\int d^3\vec{r}' \int d\omega \vec{E}(\vec{r}', \omega) \cdot \vec{j}(\vec{r}', \omega)^* \right)$$

$$\begin{aligned} \frac{d^2W}{dzd\omega} &\simeq -\operatorname{Re} \left(\frac{2i\alpha}{3\pi} \frac{\hat{q}}{E^2} \int_0^\infty d\bar{t} \frac{\omega n^2}{\epsilon} \exp \left[-\omega |n_i| \beta \bar{t} \left(1 - \frac{\hat{q}}{6E^2} \bar{t} \right) \right] \right. \\ &\quad \left. \times \cos(\omega \bar{t}) \exp \left[i\omega n_r \beta \bar{t} \left(1 - \frac{\hat{q}}{6E^2} \bar{t} \right) \right] \right) \end{aligned}$$

without damping

Introduction

QCD case

(Q)ED case


goals

arXiv:1106.2856v1

without damping

with damping

Summary



$$\Gamma = 0, \quad m = 0, 0.3, 0.6, 0.9 \text{ GeV}$$

$$\hat{q} = 2.5 \text{ GeV}^2/\text{fm}, \quad E = 20 \text{ GeV}, \quad M = 1 \text{ GeV}$$

without damping

Introduction

QCD case

(Q)ED case



goals

arXiv:1106.2856v1

without damping

with damping

Summary



$$\Gamma = 0, \quad m = 0, 0.3, 0.6, 0.9 \text{ GeV}$$

$$\hat{q} = 2.5 \text{ GeV}^2/\text{fm}, \quad E = 20 \text{ GeV}, \quad M = 1 \text{ GeV}$$

$$\frac{dW}{d\omega} = \frac{t_{\text{form}}}{t_{\text{BH}}} \times \frac{dW_{\text{BH}}}{d\omega}$$

with damping

Introduction

QCD case

(Q)ED case



goals

arXiv:1106.2856v1

without damping

with damping

Summary



$$m = 0.6 \text{ GeV}, \quad \Gamma = 0, 5, 10, 50 \text{ MeV}$$

$$\hat{q} = 2.5 \text{ GeV}^2/\text{fm}, \quad E = 20 \text{ GeV}, \quad M = 1 \text{ GeV}$$

$$\frac{dW}{d\omega} = \frac{t_{\text{form}}}{t_{\text{BH}}} \times \frac{dW_{\text{BH}}}{d\omega}$$

Summary

Introduction

QCD case

(Q)ED case

Summary

- Effect of **damping** on radiative energy loss
- Energy loss spectrum from formation time
 - light parton
 - comparison with complete calculation in ED
- Questions:
 - strength of damping in a QCD plasma?
 - visible effects in quenching?