


# Heavy-light Flavor Correlations on the QCD Phase Boundary

Chihiro Sasaki

Institute of Theoretical Physics, University of Wroclaw, Poland

- [1] C.S., Phys. Rev. D **90**, no. 11, 114007 (2014).
- [2] C.S. and K. Redlich, Phys. Rev. D **91**, no. 7, 074021 (2015).

- crossover temperatures: *not unique!*



- flavor basis vs. conserved charge basis: strange mesons deconfined at  $T_{\text{ch}}$ !

$$\mu_u = \frac{1}{3}\mu_B + \frac{2}{3}\mu_Q, \quad \mu_d = \frac{1}{3}\mu_B - \frac{1}{3}\mu_Q, \quad \mu_s = \frac{1}{3}\mu_B - \frac{1}{3}\mu_Q - \mu_S.$$

- charm? ... lessons from lattice QCD:

(i) EoS not affected by dynamical c quark around  $T_{\text{ch}}$  [Borsanyi et al. ('11)]

(ii) charmed mesons deconfined together with light mesons [Basavov et al. ('14)]

- correlations between light and heavy-flavor physics

$\Rightarrow$  how are heavy-light hadrons modified toward chiral crossover?

$$D_s \sim c\bar{s} \text{ is like } K \sim q\bar{s}? \dots \text{ NO!}$$

## **I. Chiral Structure of Heavy-light Mesons**


# Symmetries of QCD in the heavy quark mass limit

- flavor symmetries

chiral symmetry :  $m_{u,d}/\Lambda_{\text{QCD}} \ll 1$ ,  $m_s/\Lambda_{\text{QCD}} < 1$ .

heavy quark symmetry :  $\Lambda_{\text{QCD}}/m_{c,b} \ll 1$ .


- heavy-light ( $Q\bar{q}$ ) mesons  $Q$  : heavy quark and  $q$  : light quark  
e.g. D mesons:  $Q = c$ ,  $q = u, d, s$



- physical picture ( $m_Q \rightarrow \infty$ )
  - **flavor symmetry** ( $c \leftrightarrow b$ ): cloud does not feel the flavor of  $Q$ .
  - **spin symmetry**: cloud does not feel the spin of  $Q$ .

**Spin and flavor symmetries of heavy quarks are entangled!**

- $SU(2N_{Qf})$  **spin-flavor symmetry**: [Shuryak ('81), Isgur-Wise ('89)]  
light d.o.f. (q) do not feel the flavor and spin of the heavy quark (Q).



- spin partners:  $D(0^-)$  and  $D(1^-)$ ,  $B(0^-)$  and  $B(1^-)$

- **real world:**

$$m_{D^*} - m_D = 142 \text{ MeV}, m_{B^*} - m_B = 46 \text{ MeV} \ll \Lambda_{\text{QCD}}$$

...  $1/m_Q$  corrections

$$m_{D_s} - m_{D_d} = 100 \text{ MeV}, m_{B_s} - m_{B_d} = 90 \text{ MeV} \ll \Lambda_{\text{QCD}}$$

...  $m_q$  corrections

## Role of light flavor (chiral) symmetry

- **observation**: 2nd lowest spin doublets


$$D_{u,d}(0^+) : 2308 \text{ MeV} \quad [\text{Belle (03)}] \quad D_{u,d}(1^+) : 2427 \text{ MeV} \quad [\text{Belle (03)}]$$

$$D_s(0^+) : 2317 \text{ MeV} \quad [\text{Babar (03)}] \quad D_s(1^+) : 2460 \text{ MeV} \quad [\text{CLEO (03)}]$$

- mass difference of parity doublets:  $\delta m = 300 - 400 \text{ MeV} \sim \Lambda_{\text{QCD}}$

NOTE: potential model for D mesons (cf. hydrogen atom) does not work!

- chiral doubling [Nowak-Rho-Zahed (92); Bardeen-Hill (93)]



effective theory for heavy-light system based on the two relevant symmetries

## II. Thermodynamics

## Embedding $D, D_s$ in a linear sigma model

- chiral fields  $\Sigma = \sigma + i\pi$ , heavy-light meson fields  $H(0^-, 1^-), G(0^+, 1^+)$

$$\Sigma \rightarrow g_L \Sigma g_R^\dagger, \quad \mathcal{H}_{L,R} \rightarrow S \mathcal{H}_{L,R} g_{L,R}^\dagger.$$

- Lagrangian

$$\mathcal{L} = \mathcal{L}_L(\Sigma) + \mathcal{L}_{\text{HL}}(\mathcal{H}, \Sigma), \quad V_{\text{HL}} = V_{\text{HL}}(\mathcal{H}^2, \mathcal{H}^4; \Sigma) + V_{\text{HL}}^{(\text{exp})}.$$



- 6 parameters fixed with  $T = 0$  physics

$$V_{\text{HL}}^{(2)} : m_0, \underbrace{g_\pi^q, g_\pi^s}_{\Sigma \leftrightarrow \mathcal{H}^2}, \quad V_{\text{HL}}^{(4)} : k_0, \underbrace{k_q, k_s}_{\Sigma \leftrightarrow \mathcal{H}^4}$$


- isospin sym & mean field approximation:  $\langle \sigma_q \rangle, \langle \sigma_s \rangle, \langle D_q \rangle, \langle D_s \rangle$

*conventional approach ... then?*

# Chiral condensates: role of charmed-meson MF



[HotQCD Collaboration ('12)]



- lattice: qualitative diff. between  $\langle \bar{q}q \rangle$  and  $\langle \bar{s}s \rangle \dots$  **SU(2+1)**:  $T_c^{(u,d)} < T_c^{(s)}$
- chiral model:  $\sigma_{q,s}$  — approx. **SU(3)!**
- induced chiral sym. breaking:

$$h_q^* = h_q - D_q^2 \left( \frac{1}{2} g_\pi^q + 2k_q D_q^2 \right),$$

$$h_s^* = h_s - \frac{1}{\sqrt{2}} D_s^2 \left( \frac{1}{2} g_\pi^s + 2k_s D_s^2 \right).$$



*conventional approach:*

1. set up at  $T = 0$ , all the parameters are *constant*.
2. 4 gap equations at given  $T$
3. approximate SU(3)  $h_q^*/h_s^* \sim 1 \dots!?$

*resolution:*



1.  $\langle \sigma_q \rangle$  and  $\langle \sigma_s \rangle$  as input    e.g. lattice chiral condensates
2.  $\langle D_q \rangle$ ,  $\langle D_s \rangle$  and 2 HL-couplings as output     $\Rightarrow g_\pi, k$  varying with  $T$
3.  $h_q^*/h_s^* \ll 1$  restored

# Intrinsic thermal effects





- concept of EFT: generating functional, Green's functions

$$Z = \int \mathcal{D}q \mathcal{D}g e^{S_{\text{QCD}}[q,g]} \equiv \int \mathcal{D}U e^{S_{\text{eff}}[U]}$$




- low-energy constants: high-frequency modes integrated out  
 $\Rightarrow$  in a hot/dense medium: effective couplings dep. on  $T/n$
- L:  $T_{pc}^{\text{lat}} = 154 \text{ MeV} \Rightarrow$  quartic coupling such that  $m_\sigma = 400 \text{ MeV}$   
 HL:  $\sigma_{q,s}$  profiles from lattice QCD  $\Rightarrow g_\pi^{q,s}(T)$  etc.

# In-medium charmed-meson masses




- chiral splitting at  $T_{pc}$ :  $\delta M_D \simeq \delta M_{D_s}$   
*... insensitive to light flavors!*  
 $\Rightarrow$  heavy quark symmetry
- light mesons at  $T_{pc}$ :  $\delta M_{\pi-\sigma} \ll \delta M_{K-\kappa}$   
*...  $SU(2+1) \neq SU(3)$*
- cf. chiral  $SU(4)$ : [Roder-Ruppert-Rischke ('03)]  
 $\delta M_D \ll \delta M_{D_s}$



- $D_s$  screening mass from lattice QCD

[Bazavov-Karsch-Maezawa-Mukherjee-Petreczky ('14) ]



- **Quenched HL coupling and  $D_s$  decays:** anomalous suppression

$$\Gamma(0^+/1^+ \rightarrow 0^-/1^- + \pi^0) \propto \underbrace{(g_\pi^s)^2}_{\text{quenched due to CSR}} \cdot \underbrace{\delta_{\pi^0\eta}^2}_{\text{isospin violation}}$$

## Generalized susceptibilities

- generating functional vs. effective action

$$\Gamma[\phi_{\text{cl}}] = -W[J] - \int d^4x J(x)\phi_{\text{cl}}(x)$$

- fluctuation of  $\phi$

$$\langle \phi(x)\phi(y) \rangle - \langle \phi(x) \rangle \langle \phi(y) \rangle = \frac{\delta^2 W[J]}{\delta J(x)\delta J(y)} = \left( \frac{\delta^2 \Gamma[\phi]}{\delta \phi_{\text{cl}}(x)\delta \phi_{\text{cl}}(y)} \right)^{-1}$$
$$\therefore 1 = \frac{\delta^2 W}{\delta J \delta J} \frac{\delta^2 \Gamma}{\delta \phi_{\text{cl}} \delta \phi_{\text{cl}}}$$

- multiple fields  $\vec{\phi} = (\phi_1, \phi_2, \dots, \phi_n)$


$$\delta_{ij} = \frac{\delta^2 W}{\delta J_i \delta J_{\underline{k}}} \frac{\delta^2 \Gamma}{\delta \phi_{\underline{k}} \delta \phi_j}, \quad \{i, j, k\} = 1, 2, \dots, n$$

–  $2 \times 2$  sus. matrix  $\Rightarrow \chi_{qq,qs,ss} \sim \chi_{\text{ch}}$ : light flavor correlations

–  $4 \times 4$  sus. matrix  $\Rightarrow \chi_{\sigma D}, \chi_{DD}$ : heavy-light flavor correlations

# Correlations between light and heavy-light mesons

[CS-Redlich ('14)]



qualitative changes set in at  $T \sim T_{pc}$ : (NOTE:  $\chi_{ch} \sim \partial \sigma_{q,s} / \partial m_{q,s}$ )

$$\hat{\chi}_{\sigma D} = -\hat{\chi}_{ch} \hat{C}_{HL} \hat{\chi}_D, \quad \hat{\chi}_{D\sigma} = -\hat{\chi}_D \hat{C}_{HL} \hat{\chi}_{ch},$$

$$\hat{\chi}_{DD} = \hat{C}_D - \hat{C}_{HL} \hat{\chi}_{ch} \hat{C}_{HL} \equiv \hat{\chi}_D.$$

**in-medium  $D_s$  as a probe of  $O(4)$ !**

## Toward high-density QCD

- effective Lagrangian parameters *vary* with  $T$ !  
⇒ other hadrons: more and more states activated toward QCD p.t.
  - role of higher KK modes in open moose model [Son, Stephanov ('03)]  
correct high-energy behavior for current correlator
  - nuclear matter saturation in Walecka model  
density-dep. parameters: many-body effects integrated out

## How to handle them?

- holographic QCD models:  $1/N_c$  corrections?
- 4d effective theories: higher resonances, careful treatment of broad resonances ⇒ talk by Pok Man Lo
- microscopic approach: lattice QCD, DS/FRG

## Summary

- **Synthesis of light and heavy quark dynamics**

$$\frac{m_q}{m_c}, \frac{m_s}{m_c}, \frac{T}{m_c} \ll 1 \quad \text{heavy quark symmetry as a reliable guide}$$

- at  $T_{pc}$ : chiral mass splittings of HL mesons insensitive to light flavors.

$$\delta M_{D,B} \simeq \delta M_{D_s,B_s} \quad \text{vs.} \quad \delta M_{\pi-\sigma} \ll \delta M_{K-\kappa}$$

- remnant of  $O(4)$  in HL mixed fluctuations.

- anomalous suppression of  $D_s$  decay widths as a sign of CSR

**in-medium  $D_s$  as a probe of  $O(4)$ !**

- **Application to a dense system**

- strange and charm number conservation

- role of higher-lying hadrons

- chiral restoration vs. deconfinement  $\dots$  lat. Dirac-eigenmode expansion

- fluc. of conserved charges  $\chi_X^{(\text{non-reg})} = \mathcal{F}_X(\sigma_{q,s}, D_{q,s}; \chi_{ch})$