

Renormalization in Self-Consistent Approximation schemes at Finite Temperature I: Theory

Hendrik van Hees, Jörn Knoll

GSI Darmstadt

(Dated: August 29, 2001)

Within finite temperature field theory, we show that truncated non-perturbative self-consistent Dyson resummation schemes can be renormalized with local counter terms defined at the vacuum level. The requirements are that the underlying theory is renormalizable and that the self-consistent scheme follows Baym's Φ -derivable concept. The scheme generates both, the renormalized self-consistent equations of motion and the closed equations for the infinite set of counter terms. At the same time the corresponding 2PI-generating functional and the thermodynamical potential can be renormalized, in consistency with the equations of motion. This guarantees the standard Φ -derivable properties like thermodynamic consistency and exact conservation laws also for the renormalized approximation schemes to hold. The proof uses the techniques of BPHZ-renormalization to cope with the explicit and the hidden overlapping vacuum divergences.

PACS numbers: 11.10.-z, 11.10.Gh, 11.10.Wx

I. INTRODUCTION

In recent years the question how to appropriately treat particles in a hot and dense medium has continuously gained growing interest in many areas of physics ranging from plasma physics, condensed matter physics to nuclear and particles physics. Within a Green's function formalism a consistent treatment of such phenomena frequently leads to consider dressed propagators, which follow from non-perturbative Dyson resummation schemes, rather than perturbative ones, in particular, if damping width effects play a significant role.

Already in the early sixties, based on a functional formulation of Luttinger and Ward [1] and Lee and Yang [2], Kadanoff and Baym [3] considered a class of self-consistent Dyson approximations. Baym reformulated this in terms of a variational principle, defining the so called Φ -derivable approximations [4]. Since in principle the truncation of the diagrammatic series of the functional Φ can be at arbitrary level it gives rise to a variety of approximations including Hartree and Hartree-Fock as the simplest schemes. The main virtue of this concept is that the resulting equations of motion are conserving and the corresponding equilibrium limit is thermodynamically consistent. This functional treatment constitutes the basis for the two-particle irreducible (2PI) diagram technique, where the functional Φ generates the driving terms for the equations of motion, like the self-energy. Later the concept was extended to the relativistic case and formulated within the path integral approach by Cornwall, Jackiw and Tomboulis [5]. There is no formal problem to extend the formalism to the Schwinger Keldysh real-time path method [6, 7] applicable to the general case of non-equilibrium many-body theory.

Despite such early conceptual formulations most applications of self-consistent approximations were pursued on the Hartree or Hartree-Fock level sometimes supplemented by RPA resummations (see, e.g., [8, 9]) or perturbative estimates of higher order corrections. Thus, essentially mean field corrections to the self-energies were considered. Genuine two-point or even multi-point contributions to the self-consistent self-energy, which give rise to a finite damping width, imply a new level of complexity. Various new conceptual problems, like leaving the quasi-particle picture or the issue of renormalization come in with considerable complications for the numerical solutions of such problems. In the pioneering work of Bielajew and Serot [10] for the first time the renormalization of

self-consistent two-point self-energy loops were investigated at zero temperature but finite matter density.

In recent years with the special interest in dense hadronic matter problems the Φ -derivable schemes with higher order self-energy terms were used to derive transport equations [11–14] from the corresponding Kadanoff-Baym equations [15] for the consistent and also conserving [16] transport treatment of particles with finite spectral width beyond the quasi-particle approximation. Also first investigations involving finite mass width effects on vector mesons were investigated within a self-consistent Dyson resummation scheme [17]. In most of these cases, however, the question of renormalization was circumvented by taking into account the imaginary part of the self-energy only, while the real part was neglected, or cut-off recipes – mostly symmetry violating – were employed, or even the counter terms were chosen temperature dependent! Yet, especially in the study of phase transitions, e.g., within chiral hadronic models of QCD, or for non-perturbative corrections of hard thermal loop approaches to QCD, e.g., within a Φ -derivable scheme [18, 19] it is important to consistently take into account both, real and imaginary parts of the self-energies.

Therefore in this paper we address the more formal question of renormalizability of such non-perturbative approximations. We essentially concentrate on the thermodynamic equilibrium case and show how to obtain finite self-consistent dynamical quantities like the in-medium equations of motion and the self-energy of the particles and thermodynamic quantities like the pressure and the entropy. For definiteness and clarity of the presentation we use the ϕ^4 -theory as the most simple example to study the related questions. The results and techniques can easily be transferred to other theories.

The paper is organized as follows: In section II we briefly summarize Baym’s Φ -functional using the combined real and imaginary time contour appropriate for thermal equilibrium within the path integral formalism [5].

In section III we derive the general formalism for the renormalization of the self-consistent self-energy at finite temperature and the in-matter generating functional Γ . With the help of Weinberg’s convergence theorem [20] and the BPHZ-formalism of renormalization theory [21, 22] we show that, in close analogy to perturbative renormalization (see, e.g., [23–25]) *any Φ -derivable self-consistent approximation scheme can be rendered finite by subtracting pure vacuum counter terms* given by closed recursive equations. Indeed the main complication arises from the fact that the self-consistent propagator is involved in divergent loops which gives rise to “hidden” divergences which have to be resolved. This leads to a Bethe-Salpeter equation for the divergent vacuum pieces with a kernel compatible with the functional Φ , which needs to be renormalized. The renormalized equations of motion for the self-consistent propagator are shown to be consistent with the renormalized 2PI generating functional which proves the consistency of counter terms at both levels. The diagrammatical interpretation shows that in strict analogy to perturbative renormalization of thermal quantum field theory this procedure can be interpreted as renormalization of the wave functions, the mass and the coupling constants *in the vacuum*. We also give a closed expression for the renormalized self-consistent thermodynamical potential.

Numerical solutions for the renormalized self-consistent Dyson equations beyond the standard Hartree approximation up to the self-consistent sunset-diagram level could be achieved; the results are discussed in the second paper of this series [26].

II. Φ -DERIVABLE APPROXIMATIONS

In the case of thermal equilibrium the real and imaginary time formalism can be combined by extending the Schwinger-Keldysh contour $\mathcal{C}_{\mathbb{R}}$, running from t_i to t_f and back to t_i , by appending a vertical part \mathcal{C}_{Th} running from t_i to $-i\beta$ (see fig. 1). One uses the fact that the factor $\exp(-\beta\mathbf{H})$




FIG. 1: The Schwinger-Keldysh real-time contour modified for the application to thermal equilibrium of quantum field theory.

in the canonical density operator can be formally treated as a time evolution in imaginary time direction. The functional integral for bosonic fields has to be taken over all fields fulfilling the *periodic boundary condition* $\phi(t_i - i\beta) = \phi(t_i)$ which leads to the Kubo-Martin-Schwinger (KMS) condition for the Green's functions [27]. Since the equilibrium state is invariant in time one can take $t_i \rightarrow -\infty$ and $t_f \rightarrow +\infty$ which is convenient when formulating the theory in energy-momentum space via a Fourier transformation.

In addition to the usually introduced *one-point auxiliary external source* also a *two-point auxiliary external source* is included. Variations of the latter generate contour-ordered expectation values of the form $\langle \mathcal{T}_{\mathcal{C}} \phi(x_1) \phi(x_2) \rangle$ where \mathcal{C} denotes the extended Schwinger-Keldysh time contour, and $\mathcal{T}_{\mathcal{C}}$ stands for the ordering of the operators due to the ordering of the time arguments along this contour.

The corresponding generating functional is defined within the path integral formalism of quantum field theory as

$$Z[J, B] = N \int D\phi \exp \left[iS[\phi] + i \int_{\mathcal{C}} d(1) J_1 \phi_1 + \frac{i}{2} \int_{\mathcal{C}} d(12) B_{12} \phi_1 \phi_2 \right], \quad (1)$$

where N is a normalization constant chosen below. Here and in the following the shorthand notation

$$\int_{\mathcal{C}} d(12 \dots n) f_{12 \dots n} = \int_{\mathcal{C} \times \mathbb{R}^{d-1}} d^d x_1 \cdots d^d x_n f(x_1, \dots, x_n) \quad (2)$$

for integrations in d -dimensional space is used (in the sense of dimensional regularization¹). The time integration has to be performed along the time contour introduced above. It is clear that also the action functional S has to be defined as the $\mathcal{C} \times \mathbb{R}^{d-1}$ -integral of the Lagrangian.

The generating functional W for *connected diagrams* reads

$$W[J, B] = -i \ln(NZ[J, B]). \quad (3)$$

¹ We use this notation for convenient regularization only, in order to write down sensible non-renormalized functionals.

The *mean field* and the *connected Green's functions* are defined by

$$\varphi_1 = \frac{\delta W}{\delta J_1}, \quad G_{12} = -\frac{\delta^2 W}{\delta J_1 \delta J_2} \quad \Rightarrow \quad \frac{\delta W}{\delta B_{12}} = \frac{1}{2}(\varphi_1 \varphi_2 + iG_{12}). \quad (4)$$

The last formula immediately follows from the definition of the partition sum (1) and (3) using the Feynman-Kac formula

$$\langle \mathcal{I}_{\mathcal{C}} \phi(x_1) \cdots \phi(x_n) \rangle = \int D\phi \phi(x_1) \cdots \phi(x_n) \exp(iS[\phi]). \quad (5)$$

Since the real-time part of this contour is closed by itself it can be shown that the functional (1) factorizes into the real-time part and the imaginary time part. Thus the Feynman rules for the calculation of connected Green's functions apply separately to the real and the imaginary part of the contour [28], since the functional (3) splits into a sum of the contributions from the vertical and the real-time part of the contour respectively:

$$Z[J, B] = Z_{\mathbb{R}}[J, B] Z_{\text{Th}}[J, B], \quad W[J, B] = W_{\mathbb{R}}[J, B] + W_{\text{Th}}[J, B] - i \ln N. \quad (6)$$

Within the real-time part the effect of the heat bath is completely taken into account by means of the analytical properties of the Green's function. It is uniquely determined by the KMS-condition which itself is a consequence of the above mentioned periodic boundary conditions for bosons within the path integral (see appendix A for details about the analytic properties of Green's functions and self-energies).

By a functional Legendre transformation in φ and G one obtains the *effective quantum action*:

$$\Gamma[\varphi, G] = W[J, B] - \int_{\mathcal{C}} d(1) \varphi_1 J_1 - \frac{1}{2} \int_{\mathcal{C}} d(12) (\varphi_1 \varphi_2 + iG_{12}) B_{12} - i \ln N. \quad (7)$$

Now, as is well known from the usual functional formalism of quantum field theory, a formal saddle point expansion of the effective quantum action is an expansion in orders of \hbar around the classical solution, where G is considered as an independent quantity, gives

$$\Gamma[\varphi, G] = S[\varphi] + \frac{i}{2} \text{Tr} \ln(M^2 G^{-1}) + \frac{i}{2} \int_{\mathcal{C}} d(12) D_{12}^{-1} (G_{12} - D_{12}) + \Phi[\varphi, G]. \quad (8)$$

Herein the free propagator in the presence of a mean field is given by

$$D_{12}^{-1} = \frac{\delta^2 S[\varphi]}{\delta \varphi_1 \delta \varphi_2}. \quad (9)$$

The arbitrary constant M^2 account for the overall normalization and cancels for any physical quantities as we shall see below.

In the case of an ideal gas it is sufficient to subtract the pure vacuum part to render this functional finite, which leads to the well known result. At $T = 0$ this subtraction corresponds to the renormalization of the total ground-state energy to zero.

As we shall discuss below in the case of interacting particles this description is not sufficient to render the effective action and thus the pressure finite, since we need additional subtractions of vacuum sub-divergences to renormalize it.

In the above sense the functional Φ in (8) contains the parts of order $O(\hbar^2)$ and higher. Since simple power counting shows the \hbar -order of diagrams to be identical with the number of loops, Φ as a functional of G consists of all closed diagrams with at least 2 loops. The lines within the diagrams stand for dressed Green's functions G while the vertices are the bare vertices of the classical action

with presence of the background field φ which can be immediately read off from $S_I[\varphi + \phi']$ around $\phi' = 0$ beginning at order ϕ'^3 .

The equations of motion are now given by the fact that we like to study the theory with vanishing auxiliary sources j and B . From (4) and (7) we obtain

$$\frac{\delta\Gamma}{\delta\varphi_1} = -j_1 \stackrel{!}{=} 0, \quad \frac{\delta\Gamma}{\delta G_{12}} = -\frac{i}{2}B_{12} \stackrel{!}{=} 0. \quad (10)$$

Using (8) the equations of motion read

$$\begin{aligned} \frac{\delta S}{\delta\varphi_1} &= -\frac{i}{2} \int_{\mathcal{C}} d(1'2') \frac{\delta D_{1'2'}^{-1}}{\delta\varphi_1} G_{1'2'} - \frac{\delta\Phi}{\delta\varphi_1} \\ \Sigma_{12} &:= D_{12}^{-1} - G_{12}^{-1} = 2i \frac{\delta\Phi}{\delta G_{12}}. \end{aligned} \quad (11)$$

The first line is of the form of a Klein-Gordon equation with the quantum corrections to the classical theory on the right hand side. The second equation is the *Dyson equation* and shows that the variation of the Φ -functional with respect to G is the self-energy defined with respect to the classical field dependent propagator (9). This shows that the Φ -functional must be *two-particle irreducible*. No propagator line must contain a self-energy insertion. In other words the closed diagrams representing contributions to Φ must not split in disconnected pieces when cutting two lines. Diagrammatically the derivative of a functional with respect to G corresponds to opening one line of the diagrams representing it. In that sense Φ is the generating functional for *skeleton diagrams for the self-energy* where the lines represent fully dressed propagators. Thus the functional formalism avoids double counting in a natural way by omitting all non-skeleton diagrams from the Dyson-resummed equations of motion. Altogether the Φ -functional formalism provides a closed system of exact equations of motion for the full 2-point function and the full mean field. Solving these equations would be equivalent to finding the full propagator of the quantum field theory which of course is impossible in practice.

One obtains approximations by truncating the series for Φ at a certain vertex or loop order (which corresponds to the respective order in the coupling λ or \hbar respectively), while preserving the forms (10) and (11) of the self-consistent equations of motion. Approximations of this kind respect the conservation laws for the expectation values of Noether currents for symmetries which are linearly operating on the field operators (including space-time symmetries and the according conserved quantities as energy, momentum and angular momentum) [4, 11].

In the case of thermal equilibrium, setting the mean field and the propagator to the solution of the self-consistent equations the effective action gives the grand canonical potential $\Omega = -T\Gamma[\varphi, G]_{J,K=0} = -T \ln Z(\beta)$ [27]. Since the real-time part of the contour in figure 1 is closed it vanishes for the solution of the equations of motion. A short summary about the analytic continuation from the real to the imaginary time formalism is given in appendix A.

Thus the formalism leads to a well defined treatment for bulk thermodynamical quantities of the system (like energy, pressure, entropy, etc.).

All these quantities can be calculated either with real-time Green's functions or with the corresponding imaginary time functions, because as summarized in appendix A real and imaginary-time propagators are connected by the analytic properties of the Green's functions originating from the KMS-condition. For our purpose the real-time formalism is preferred, because of its simplicity with respect to the analytic structure of Green's functions, which easily permits to deal with the mixture of finite temperature and vacuum pieces occurring in the subtraction scheme. This also avoids the necessity to perform an analytic continuation from imaginary time to real-time Green's functions which is complicated to obtain for numerical results.

In order to exemplify the method we apply the formalism to ϕ^4 -theory with the Lagrangian

$$\mathcal{L} = \frac{1}{2}(\partial_\mu\phi)(\partial^\mu\phi) - \frac{m^2}{2}\phi^2 - \frac{\lambda}{4!}\phi^4. \quad (12)$$

III. RENORMALIZABILITY OF Φ -DERIVABLE APPROXIMATIONS

In this section we show that in close analogy to the renormalization of perturbative diagrams also any Φ -derivable self-consistent approximation can be renormalized with help of local temperature-independent counter terms.

The proof uses the same line of arguments as in the perturbative case: The reason is, that the renormalization theory completely rests on Weinberg's power counting theorem [20] which is formulated for a general class of Green's functions with a given asymptotic behavior. It does not depend on the special form of the propagators.

The first step is a simple topological argument leading to the superficial degree of divergence for a given diagram γ , which for the ϕ^4 theory simply follows from the number E of external lines [29]

$$\delta(\gamma) = 4 - E. \quad (13)$$

Due to field reflection symmetry only diagrams with an even number of external lines are different from 0, the only divergent functions are those represented by diagrams where the number of external legs is 0 (i.e., contributions to the total mean energy and the thermodynamical potential), 2 (self-energy (Green's function)) and 4 (four-point vertex functions).

The second step is an expansion of the regularized un-renormalized self-consistent self-energy around the self-consistent vacuum propagator which shows that the asymptotic behavior of the diagrams and sub-diagrams is ruled by their pure vacuum parts.

Then an equation of motion for the temperature-dependent "infinite part" of the regularized self-energy is derived and it is shown that it can be renormalized by a temperature independent subtraction procedure.

A. BPHZ-Scheme for the vacuum

We first apply the BPHZ renormalization theorem [21, 22, 30] for the vacuum. The only difference to the perturbative case is that we apply it to diagrams with self-consistent propagator lines. This is justified since Weinberg's power counting theorem is independent of the special form of propagators but only needs their asymptotic behavior stated above.

We summarize the BPHZ scheme as follows. A sub-diagram γ of a diagram Γ is defined as any set of lines and vertices contained in Γ which itself builds a proper vertex diagram: $\gamma \subseteq \Gamma$. A sub-diagram γ is called *renormalization part* if its superficial degree of divergence (or its dimension) is greater than or equal to 0. In our case this means it has at most four external legs. Two sub-diagrams γ_1 and γ_2 are called *nested* $\gamma_1 \subseteq \gamma_2$ if γ_1 is a sub-diagram of γ_2 . If they have no line or vertex in common, $\gamma_1 \cap \gamma_2 = \emptyset$, they are called *disjoined*. If they are neither nested nor disjoined they are called *overlapping*: $\gamma_1 \circ \gamma_2$.

To any diagram Γ we denote the integrand following from the Feynman rules with I_Γ . For a set of pairwise disjoined sub-diagrams $\gamma_1, \gamma_2, \dots, \gamma_n$ we write the integrand in terms of the integrands of the sub-diagrams I_{γ_j} and the remainder of the integrand denoted by $I_{\Gamma \setminus \{\gamma_1, \dots, \gamma_n\}}$, usually called

the *reduced diagram*:

$$I_\Gamma = I_{\Gamma \setminus \{\gamma_1, \dots, \gamma_n\}} \prod_{j=1}^n I_{\gamma_j}. \quad (14)$$

The original scheme by Bogoliubov and Parasiuk [21] defines recursively the integrand R_Γ of the renormalized diagram. If a diagram does not contain any renormalization part but is itself divergent it is called *primitively* divergent. In that case the renormalized integrand is defined by $R_\Gamma = (1 - t_\Gamma)I_\Gamma$. Herein t_Γ is the operator of the Taylor expansion with respect to the external momenta around 0 up to the order of the dimension $\delta(\gamma)$ of the divergent diagram, which is in our case $4 - E$:

$$t_\gamma I_\gamma(p_1, \dots, p_k) := \begin{cases} \sum_{j=0}^{\delta(\gamma)} \frac{1}{j!} \sum_{\substack{\mu_1, \dots, \mu_k \geq 0 \\ \mu_1 + \dots + \mu_k = j}} \frac{\partial^j I_\gamma(p_1, \dots, p_k)}{\partial p_1^{\mu_1} \dots \partial p_k^{\mu_k}} \Big|_{p_1 = \dots = p_k = 0} p_1^{\mu_1} \dots p_k^{\mu_k} \\ 0 \quad \text{for} \quad \delta(\gamma) < 0 \end{cases} \quad (15)$$

If the diagram is convergent the integrand is unchanged under renormalization.

If the diagram is not only primitively divergent but contains divergent sub-diagrams the integrand for the diagram with all subdivergences subtracted is called \bar{R}_Γ and the renormalized integrand is defined by

$$R_\Gamma = \begin{cases} \bar{R}_\Gamma & \text{if } \delta(\Gamma) < 0 \\ (1 - t_\Gamma)\bar{R}_\Gamma & \text{if } \delta(\Gamma) \geq 0. \end{cases} \quad (16)$$

From Weinberg's power counting theorem it follows that after this recursive procedure the integral over the internal momenta of R_Γ is finite. The definition of the *counter terms* by the Taylor operator t_γ for any renormalization part γ of the diagram shows that these are polynomials in the external momenta to the order $\delta(\gamma)$ and thus can be interpreted as counter terms to the corresponding wave function normalization factors, masses and coupling constants in the original Lagrangian.

Zimmermann solved Bogoliubov's and Parasiuk's recursion with his *forest* formula. A forest is defined as any set of sub-diagrams (including the empty set and the whole diagram itself) which are *pairwise non-overlapping*. One can depict these sets by drawing boxes around the sub-diagrams and in a forest these boxes are not allowed to overlap but they can be nested. A forest is restricted if each of its boxes contains only renormalization parts. To each restricted forest \mathfrak{F} one associates again an integrand, namely

$$\Omega_{\mathfrak{F}} = \prod_{\gamma \in \mathfrak{F}} (-t_\gamma) I_\gamma = \begin{array}{c} \boxed{\text{---}i\Gamma^{(4)}\text{---}} \\ \text{---} \\ \boxed{\text{---}i\Gamma^{(4)}\text{---}} \\ \text{---} \\ \text{---} \\ \boxed{\text{---}i\Gamma^{(4)}\text{---}} \\ \text{---} \\ \text{---} \\ \boxed{\text{---}i\Gamma^{(4)}\text{---}} \\ \text{---} \\ \text{---} \\ \boxed{\text{---}i\Gamma^{(4)}\text{---}} \end{array} \quad (17)$$

The diagram to the right shows an example case for a typical ladder diagram, which we shall consider in the following section. For this case $\delta(\gamma) = 0$ for all sub-diagrams and the diagram itself such that for each box only the subdiagram value at vanishing external momenta is to be subtracted. The tilde over the product sign in (17) stands for the fact that in case of nested diagrams within the forest one has to apply the Taylor operators from the innermost to the outermost diagrams while for disjointed sub-diagrams the expressions are naturally independent of the order of Taylor operators, since then

$$I_\Gamma = I_{\Gamma \setminus \{\gamma_1, \dots, \gamma_n\}} \prod_{k=1}^n I_{\gamma_k}. \quad (18)$$

The forest formula then says that the integrand of the renormalized diagram is given by the sum over all restricted forests:

$$R_\Gamma = \sum_{\mathfrak{F} \in \mathcal{F}_R(\Gamma)} \Omega_{\mathfrak{F}}. \quad (19)$$

It is understood that the empty set stands for the diagram itself, i.e., without any box around a sub-diagram.

The described BPHZ-scheme chooses the renormalization point for the divergent diagrams at external momenta set to 0. It is clear that by another finite renormalization we can switch to any renormalization scheme appropriate for the application considered. In our case of ϕ^4 -theory we chose the on-shell scheme. We have to define the coupling constant, the mass and the wave function normalization. This can be formulated in terms of the proper self-energy and the proper four-point vertex (the three-point vertex can be set identical to 0 because of symmetry under field reflection $\phi \rightarrow -\phi$ without destroying the renormalizability of the theory, so that we do not have to consider terms linear or cubic in the fields within the Lagrangian):

$$\Gamma^{(4, \text{vac})}(s, t, u = 0) = \frac{\lambda}{2}, \quad \Sigma^{(\text{vac})}(p^2 = m^2) = 0, \quad \partial_{p^2} \Sigma^{(\text{vac})}(p^2 = m^2) = 0. \quad (20)$$

Here s, t, u are the usual Mandelstam variables for two particle scattering, p is the external momentum of the self-energy and m^2 is the renormalized mass of the particles. The first condition defines the coupling constant at vanishing momentum transfer for the two-particle scattering to be given by λ , the second condition chooses m to be the physical mass of the particles, while the third condition ensures that the residuum of the propagator at $p^2 = m^2$ is 1 and thus the on-shell wave function is normalized to 1 as it should be.


B. The finite temperature self-energy at the regularized level

In this section we like to isolate those vacuum subparts inherent in the pure temperature part of the self-energy which need to be renormalized at physical space-time dimension $d = 4$. For this purpose we assume a regularization scheme, e.g., dimensional regularization and extract those vacuum parts from the self-energy which diverge in the limit $d \rightarrow 4$. For sake of clarification we mark all equations with an asterisk which diverge in the limit $d \rightarrow 4$ and which need special renormalization treatment. All other equations are generally valid, even if all expressions are replaced by their renormalized quantities.

In order to extract the divergent vacuum pieces we take the self-energy as functional of the self-consistent propagator and expand it around the vacuum value

$$\Sigma_{12} = \Sigma_{12}^{(\text{vac})} + \underbrace{\Sigma_{12}^{(0)} + \Sigma_{12}^{(r)}}_{\Sigma_{12}^{(\text{matter})}} \quad (21)$$

Here $\Sigma_{12}^{(\text{vac})}$ is the vacuum ($T = 0$) self-energy. Its renormalization poses no particular problem and can be done according to standard rules. At the examples discussed in our second paper [26] it is shown how to do this in practical cases for the numerical solutions to the self-consistent equations of motion. The second and third terms in (21) contain the in-matter or finite temperature components of the self-energy. Thereby

$$-i\Sigma_{12}^{(0)} = \int_{\mathcal{C}} d(1'2') \left(-\frac{\delta i\Sigma_{12}}{\delta G_{1'2'}} \Big|_{T=0} G_{1'2'}^{(\text{matter})} \right) = \text{Diagram} \quad (22^*)$$


contains the parts of Σ linear in the matter (temperature) part $G^{(\text{matter})}$ of the full propagator

$$\begin{aligned} iG &= iG^{(\text{vac})} + iG^{(\text{matter})} \\ \underline{\underline{iG}} &= \underline{iG^{(\text{vac})}} + \text{wavy line} \end{aligned} \quad (23)$$

At this level it is important to recognize that all loops involving vertices from both sides of the real-time contour (cf. Fig. 1) are UV convergent due to the analytical properties of $\{-+\}$ and $\{+-\}$ -propagators (A16) to (A18). Such loops contain at least one thermal weight factor $n(p_0)$ which decays exponentially at large $|p_0|$, since the Θ -function parts completely drop out for large loop momenta. Therefore all mixed components like G^{-+} , or Σ^{+-} have to be excluded from the subtraction scheme. Thus, the expansion point, $G^{(\text{vac})}$, in (23) is defined as the contour-diagonal part of the propagator in the vacuum limit ($T \rightarrow 0$), i.e., with vanishing $G^{-+(\text{vac})} = G^{+-(\text{vac})} := 0$. Likewise all vacuum structures like $\Sigma^{(\text{vac})}$, and the four-point functions $\Gamma^{(4,\text{vac})}$ and $\Lambda^{(\text{vac})}$ defined below are “diagonal” in the real-time contour placement.

The remaining self-energy piece $\Sigma^{(r)}$ in (21) contains at least two $G^{(\text{matter})}$ lines which therefore are never involved in any divergent loops due to the 2PI property of the Φ -functional². Thus, there are no hidden subdivergences in $\Sigma^{(r)}$, and possible divergent vacuum sub-structures can directly be renormalized using the BPHZ rules given above.

On the other hand the diagrams of $\Sigma^{(0)}$ deserve special attention, since there the single $G^{(\text{matter})}$ -line is involved in logarithmically divergent loops, if all vertices of $\Gamma_{12,1'2'}^{(4)}$ are placed on the same side of the contour. As mentioned the terms with mixed vertices are finite.

The divergences result from the fact that the functional variation of Σ with respect to G at $T = 0$ defines a vacuum vertex function

$$-i\Gamma_{12,1'2'}^{(4)} = -\frac{\delta \Sigma_{12}}{\delta G_{1'2'}} \Big|_{T=0} = -2i \frac{\delta^2 \Phi}{\delta G_{12} \delta G_{1'2'}} \Big|_{T=0} \quad (24)$$

with four external legs.

Its diagonal part (all vertices on one contour side) defines $\Gamma_{12,1'2'}^{(4,\text{vac})}$ which is of divergence degree 0. Assuming $G^{(\text{matter})}$ of divergence degree -4 it is involved in a logarithmically divergent loop. Thus, this part of $\Sigma^{(0)}$, called $\Sigma^{(0,\text{div})}$, accounts for all terms of divergence degree 0 and consequently $\Sigma^{(r)}$ is of divergence degree -2 .

In order to trace all subdivergences hidden in $\Sigma^{(0)}$, the vacuum structure inherent in $G^{(\text{matter})}$ has to be resolved. For this purpose the diagonal parts of the full propagator (i.e., G^{--} and G^{++})

² Any $G^{(\text{matter})}$ -line in $\Sigma^{(r)}$ is either involved in loops with further $G^{(\text{matter})}$ -lines which are finite or it is attached to a pure vacuum piece. Due to the 2PI property of Φ this vacuum piece has more than four external legs also leading to finite loops for this $G^{(\text{matter})}$ -line.

renormalized, the self-energy is renormalized too, within a temperature independent subtraction scheme.

Since Φ is 2PI, the variational relation (24) defines the BS-kernel $\Gamma_{12,1'2'}^{(4,\text{vac})}$ as a proper skeleton diagram, i.e., it contains no self-energy insertions and cutting the diagram such that the pairs of space time points (12) and (1'2') are separated, cuts more than two lines. Thus the BS-kernel has the appropriate irreducibility properties for the resummation to the complete four-point function $\Lambda^{(\text{vac})}$, again showing the virtue of the Φ -functional formalism to avoid double counting.

From (28) it is obvious that switching to the momentum space representation both $\Gamma^{(4,\text{vac})}$ and $\Lambda^{(\text{vac})}$ are not needed in their full momentum dependence but rather only as a function of the two momenta given by the Fourier transformation with respect to the space-time point pairs (12) and (1'2'). Due to (24) and through (29*) both, $\Gamma^{(4,\text{vac})}$ and $\Lambda^{(\text{vac})}$, obey the symmetry relations

$$\begin{aligned} \Gamma_{12,1'2'}^{(4,\text{vac})} &= \Gamma_{1'2',12}^{(4,\text{vac})} & \text{or} & & \Gamma^{(4,\text{vac})}(p, q) &= \Gamma^{(4,\text{vac})}(q, p), \\ \Lambda_{12,1'2'}^{(\text{vac})} &= \Lambda_{1'2',12}^{(\text{vac})} & \text{or} & & \Lambda^{(\text{vac})}(p, q) &= \Lambda^{(\text{vac})}(q, p). \end{aligned} \quad (31)$$

C. Renormalization of the vacuum Bethe-Salpeter equation


In energy-momentum representation the regularized BS-equations (29*) and its equivalent “adjoint” version become

$$\Lambda^{(\text{vac})}(p, q) = \Gamma^{(4,\text{vac})}(p, q) + i \int \frac{d^d l}{(2\pi)^d} \Gamma^{(4,\text{vac})}(p, l) [G^{(\text{vac})}(l)]^2 \Lambda^{(\text{vac})}(l, q) \quad (32^*)$$

$$= \Gamma^{(4,\text{vac})}(p, q) + i \int \frac{d^d l}{(2\pi)^d} \Lambda^{(\text{vac})}(p, l) [G^{(\text{vac})}(l)]^2 \Gamma^{(4,\text{vac})}(l, q). \quad (33^*)$$

The renormalization of the BS-equations (29*) is not straight forward. First the BS-kernel $\Gamma^{(4,\text{vac})}(p, q)$ has to be renormalized following the BPHZ-rules outlined in sect. III. Representing $\Lambda(p, q)$ as the sum of ladder diagrams, this BS-kernel forms the rungs in each ladder. The complication arises from overlapping sub-divergences: in the general case each internal rung is part of various diverging sub-diagrams through the loops involving two or more rungs. However, two observations help to settle this renormalization issue:

- (i) The rungs given by the BS-kernel are 2PI with respect to cuts separating the top from the bottom extremities of the rung. This implies that there are no divergent sub-diagrams which cut into the inner structure of any rung, since this would involve a cut of more than two lines and the resulting sub-diagram would have more than four external lines. Thus divergent sub-diagrams always have complete rungs as sub-diagrams, as shown to the right.
- (ii) If one takes the difference of two Λ functions which differ only in one of the momenta, e.g., $\Lambda(p, q) - \Lambda(p', q)$, all those counter terms cancel out which contain boxes which cut the outer p or p' lines, respectively, since for these counter terms the argument p or p' are replaced by zero. Thus, the only boxes which are left are those which exclude the outermost rung attached to the p -lines. These boxes, however, just define the renormalized result of the sub-diagram complementary to this outermost rung, which is again a ladder diagram.



(valid counter term)

This permits to establish recursive relations for the renormalized expressions of the two possible differences

$$\Lambda^{(\text{ren})}(p, q) - \Lambda^{(\text{ren})}(p', q)\Gamma^{(4, \text{vac})}(p, q) - \Gamma^{(4, \text{vac})}(p', q) \quad (34)$$

$$+ i \int \frac{d^4 l}{(2\pi)^4} [\Gamma^{(4, \text{vac})}(p, l) - \Gamma^{(4, \text{vac})}(p', l)] [G^{(\text{vac})}(l)]^2 \Lambda^{(\text{ren})}(l, q),$$

$$\Lambda^{(\text{ren})}(p', q) - \Lambda^{(\text{ren})}(p', q')\Gamma^{(4, \text{vac})}(p', q) - \Gamma^{(4, \text{vac})}(p', q') \quad (35)$$

$$+ i \int \frac{d^4 l}{(2\pi)^4} \Lambda^{(\text{ren})}(p', l) [G^{(\text{vac})}(l)]^2 [\Gamma^{(4, \text{vac})}(l, q) - \Gamma^{(4, \text{vac})}(l, q')],$$

where now $\Gamma^{(4, \text{vac})}$ stands for the renormalized Bethe-Salpeter kernel. Since the renormalized function $\Gamma^{(4, \text{vac})}(p, l) - \Gamma^{(4, \text{vac})}(0, l)$ is of divergence degree less than zero the integrals are finite. This set of renormalized equations can be used to construct the renormalized Λ -function using

$$\Lambda^{(\text{ren})}(0, 0) := \pm \frac{\lambda}{2} \quad (36)$$

on the two real-time branches \mathcal{C}_{\mp} due to our renormalization condition (20). In a kind of sweep-up sweep-down scheme first the “half sided” $\Lambda^{(\text{ren})}(0, q)$ function can be constructed by solving (35) for $p' = 0$. Using this half sided function as the input for (34) the full momentum dependence of $\Lambda^{(\text{ren})}(p, q)$ can be obtained. This scheme fully complies with the BPHZ renormalization prescription. It has the remarkable feature that, although it is explicitly asymmetric in p and q , it constructs a completely symmetric renormalized four-point function which can be combined to the complete result

$$\Lambda^{(\text{ren})}(p, q) = \Gamma^{(4, \text{vac})}(p, q) + i \int \frac{d^4 l}{(2\pi)^4} [\Gamma^{(4, \text{vac})}(p, l) - \Gamma^{(4, \text{vac})}(0, l)] [G^{(\text{vac})}(l)]^2 \Lambda^{(\text{vac})}(l, q) + i \int \frac{d^4 l}{(2\pi)^4} \Lambda^{(\text{ren})}(0, l) [G^{(\text{vac})}(l)]^2 [\Gamma^{(4, \text{vac})}(l, q) - \Gamma^{(4, \text{vac})}(l, 0)]. \quad (37)$$

For numerical applications it is important to realize that only the half sided $\Lambda^{(\text{ren})}(0, l)$, and not the full momentum dependence $\Lambda^{(\text{ren})}(p, q)$, is explicitly needed. Since the half side Λ has essentially two-point function properties it can be numerically constructed using similar techniques as for self-energies.

Indeed one can express the complete renormalized self-energy part linear in $G^{(\text{matter})}$ in the form

$$\begin{aligned} \Sigma^{(0)}(p) &= \Sigma^{(0)}(p) - \Sigma^{(0)}(0) + \Sigma^{(0)}(0) \\ &= \int \frac{d^4 l}{(2\pi)^4} [\Gamma^{(4)}(p, l) - \Gamma^{(4, \text{vac})}(0, l)] G^{(\text{matter})}(l) \\ &\quad + \int \frac{d^4 l}{(2\pi)^4} \Lambda^{(\text{ren})}(0, l) G^{(\text{r})}(l) \end{aligned} \quad (38)$$

$=$

with $G^{(\text{matter})}$ and $G^{(\text{r})}$ from (23) and (25). Here $\Gamma^{(4)}$ is the full contour valued kernel (24) including mixed contour vertices, while $\Gamma^{(4,\text{vac})}$ and $\Lambda^{(\text{ren})}$ are diagonal in the contour vertices. Due to the 2PI-properties of $\Gamma^{(4)}$ the difference $\Gamma^{(4)}(l, p) - \Gamma^{(4,\text{vac})}(l, 0)$ is of divergence degree less than 0. Therefore the first integral represented by the difference of the first two diagrams is finite, since $G^{(\text{matter})}$ is of divergence degree -4 . This difference represents the most naive subtraction, which by itself, however, would be false, since it contains temperature dependent counter terms. The heart of the above derivation is that these false T -dependent counter terms are precisely compensated by the last term. The fact that the counter term structures never mix the two real-time contour branches also lifts the problem of pinch singularities which otherwise could arise due to the vanishing external momentum.

This completes the proof that the self-energies can be renormalized with T -independent counter terms.

D. Renormalization of the real-time Γ -functional

In this section we derive the renormalized real-time generating functional for the in-matter equations of motion. Thus, we restrict the contour integrations to the real-time contour $\mathcal{C}_{\mathbb{R}}$ and the corresponding traces to the real-time traces $\text{Tr}_{\mathbb{R}}$. For functions in momentum representation the corresponding contour matrix algebra in the $\{-+\}$ notation (cf. Appendix A) is implied.

For the renormalization procedure we use the ansatz (25) for the full propagator G together with the form (28) for the logarithmically divergent part of the self-energy, where the renormalized four-point function Λ resolves the subdivergences hidden in both, the propagator and the self-energy. For this purpose we decompose the generating functional Γ in its vacuum part, which is solely a functional of $G^{(\text{vac})}$, and the in-matter part

$$\Gamma = \Gamma^{(\text{vac})}[G^{(\text{vac})}] + \Gamma^{(\text{matter})}[G^{(\text{vac})}, G^{(\text{matter})}] \quad (39)$$

Thereby it is implied that the vacuum problem is already solved through its equation of motion resulting from the functional variation of $\Gamma^{(\text{vac})}$. Given $G^{(\text{vac})}$ the equations of motion in matter result from the functional variation of $\Gamma^{(\text{matter})}$ with respect to $G^{(\text{matter})}$.

Compared to the two-point self-energy the Γ - and Φ -functionals have no external points and essentially result from the diagrams of the self-energies by closing the extremities. Therefore one has to explicitly expand the corresponding expressions up to second order in $G^{(\text{matter})}$ before one comes to the situation where the remaining pieces are void of hidden subdivergences. Thus we write

$$\begin{aligned} \Phi_{\mathbb{R}} &= \Phi_{\mathbb{R}}^{(\text{vac})} + \text{Tr}_{\mathbb{R}} \frac{\delta \Phi^{(\text{vac})}}{\delta G^{(\text{vac})}} G^{(\text{matter})} + \frac{1}{2!} G^{(\text{matter})} \frac{\delta^2 \Phi^{(\text{vac})}}{\delta G^{(\text{vac})^2}} G^{(\text{matter})} + \Phi_{\mathbb{R}}^{(\text{r})} \\ &= \Phi_{\mathbb{R}}^{(\text{vac})} - \frac{i}{2} \text{Tr}_{\mathbb{R}} G^{(\text{matter})} \Sigma^{(\text{vac})} + \frac{1}{4} G^{(\text{matter})} \Gamma^{(4)} G^{(\text{matter})} + \Phi_{\mathbb{R}}^{(\text{r})}. \end{aligned} \quad (40)$$

Here we have used (11) and (24) for the vacuum parts defined through the variation of Φ with respect to G . At the same time we introduced the real-time trace in momentum space

$$\text{Tr}_{\mathbb{R}} A \cdots B = \int \frac{d^d l}{(2\pi)^d} A(l) \cdots B(l) \quad (41)$$

and the functional tensor contraction for four-point functions with propagators:

$$G_1 \Gamma^{(4)} G_2 = \int \frac{d^d l_1}{(2\pi)^d} \int \frac{d^d l_2}{(2\pi)^d} G_1(l_1) \Gamma^{(4)}(l_1, l_2) G_2(l_2). \quad (42)$$

In all expressions the functions are contour matrix functions which imply the corresponding contour matrix algebra and the contour trace, cf. Eqs. (A19) and (A20).

Applying the arguments given for $\Sigma^{(r)}$ also for $\Phi^{(r)}$ no $G^{(\text{matter})}$ -line is involved in divergent loops such that after renormalization of possible vacuum sub-divergences the entire diagram is finite. Thus only terms with at most two $G^{(\text{matter})}$ -lines need further care.

Since Φ by itself is not an observable we directly step to the definition of the Γ -functional which relates to the thermodynamic potential in the equilibrium case. Exploiting the stationarity condition of Γ at the vacuum level, i.e., using the vacuum equations of motion for $G^{(\text{vac})}$, all terms linear in $G^{(\text{matter})}$ drop out and we find for the functional with all proper vacuum sub-divergences subtracted

$$\begin{aligned} \bar{\Gamma}_{\mathbb{R}}^{(\text{matter})}[G^{(\text{matter})}] &= \frac{i}{2} \text{Tr}_{\mathbb{R}} \left(G^{(\text{matter})} \Sigma^{(\text{matter})} - \sum_{k=2}^{\infty} \frac{(G^{(\text{vac})} \Sigma^{(\text{matter})})^k}{k} \right) \\ &\quad - \frac{i}{4} \text{Tr}_{\mathbb{R}} G^{(\text{matter})} \Sigma^{(0)} + \bar{\Phi}_{\mathbb{R}}^{(r)}. \end{aligned} \quad (43^*)$$

Renormalized to zero at the vacuum level this expression is a functional of the in-matter part of the propagator. The remaining divergent parts of $\bar{\Gamma}^{(\text{matter})}$ arise from terms quadratic in $G^{(\text{matter})}$, i.e.,

$$\bar{\Gamma}_{\mathbb{R}}^{(\text{matter,div})} = \frac{i}{4} \text{Tr}_{\mathbb{R}} G^{(0)} \Sigma^{(0)} + \frac{1}{4} G^{(0)} \Gamma^{(4,\text{vac})} G^{(0)} + \frac{1}{2} G^{(0)} \Gamma^{(4,\text{vac})} G^{(r)} \quad (44^*)$$

with

$$G^{(0)} = G^{(\text{vac})} \Sigma^{(0,\text{div})} G^{(\text{vac})}, \quad (45)$$

where again in both relations above all expressions are contour diagonal. Both, $\Sigma^{(0,\text{div})}$ and $G^{(0)}$ are linear in $G^{(\text{matter})}$. Using the equations of motion for $\Sigma^{(0,\text{div})}$ and $\Lambda^{(\text{vac})}$ one arrives at an expression for the divergent part of $\bar{\Gamma}^{(\text{matter})}$ which only contains quantities which were already renormalized in the previous subsection

$$\Gamma_{\mathbb{R}}^{(\text{matter,div,ren})} = \frac{1}{4} \left(G^{(r)} \Lambda^{(\text{ren})} G^{(r)} - G^{(r)} \Gamma^{(4,\text{vac})} G^{(r)} \right). \quad (46)$$

Substituting this for the divergent part we obtain after some algebraic simplifications

$$\begin{aligned} \Gamma_{\mathbb{R}}^{(\text{matter,ren})}[G^{(\text{matter})}] &= \frac{i}{2} \overline{\text{Tr}}_{\mathbb{R}} \left(G^{(r)} \Sigma^{(\text{matter})} - \frac{1}{2} G^{(\text{vac})} \Sigma^{(r)} G^{(\text{vac})} \Sigma^{(r)} \right. \\ &\quad \left. - \sum_{k=3}^{\infty} \frac{(G^{(\text{vac})} \Sigma^{(\text{matter})})^k}{k} \right) + \frac{1}{4} G^{(r)} \Lambda^{(\text{vac})} G^{(r)} + \Phi_{\mathbb{R}}^{(r)} \\ &\quad + \text{mixed contour Tr}_{\mathbb{R}}\text{-terms from (43*)}, \end{aligned} \quad (47)$$

where $\overline{\text{Tr}}_{\mathbb{R}}$ includes only the contour diagonal parts. This expression, which now can be considered as a functional of $G^{(\text{matter})}$, or through Eqs. (25) and (28) of $G^{(r)}$, is void of any hidden subdivergences, since all matter or T -dependent parts of the propagator like $G^{(\text{matter})}$ or $G^{(r)}$ are involved in convergent loops.

Now it remains to be proven that this renormalization procedure for the Γ -functional is consistent with the renormalization of the self-energy given in the previous section. In other words: We like to show that the vanishing functional variation of $\Gamma^{(\text{ren})}[G^{(\text{matter})}]$ complies with the Dyson equation of motion and the renormalized self-energy. From the BPHZ-formalism we expect this

to hold true, because it ensures that sub-divergences can be renormalized first and then the remaining divergences which come into the game by closing the diagrams: The renormalized result is independent of the order of counter term subtractions.

It is sufficient to show this for the contour diagonal parts of (47). Writing the functional variation of $\Gamma^{(\text{ren})}$ as

$$\delta\Gamma_{\mathbb{R}}^{(\text{matter,ren})} = \frac{\delta\Gamma_{\mathbb{R}}^{(\text{matter,ren})}}{\delta\Sigma^{(r)}}\delta\Sigma^{(r)} + \frac{\delta\Gamma_{\mathbb{R}}^{(\text{matter,ren})}}{\delta G^{(r)}}\delta G^{(r)}, \quad (48)$$

where $\Sigma^{(r)}$ is supposed to be a functional of $G^{(r)}$. Both terms in (48) independently vanish. The first term drops by virtue of the Dyson equation which together with (25) ensures

$$\delta\left(\sum_{k=3}^{\infty} \frac{(G^{(\text{vac})}\Sigma^{(\text{matter})})^k}{k}\right) = \left(G^{(r)} - G^{(\text{vac})}\Sigma^{(r)}G^{(\text{vac})}\right)\delta\Sigma^{(\text{matter})}. \quad (49)$$

The second term shrinks to

$$\delta\Gamma_{\mathbb{R}}^{(\text{matter,ren})} = \frac{i}{2} \text{Tr}_{\mathbb{R}} \underbrace{\left(\Sigma^{(r)}\delta G^{(r)} + G^{(\text{vac})}\Sigma^{(r)}G^{(\text{vac})}\delta\Sigma^{(0)}\right)}_{\Sigma^{(r)}\delta G^{(\text{matter})}} + \delta\Phi_{\mathbb{R}}^{(r)} \stackrel{!}{=} 0, \quad (50)$$

which indeed implies

$$\Sigma^{(r)} = 2i \frac{\delta\Phi_{\mathbb{R}}^{(r)}}{\delta G^{(\text{matter})}}, \quad (51)$$

compatible with the definition of $\Sigma^{(r)}$. It is important to note that through the functional variation (48) only convergent loops are opened, such that none of the counter terms is affected by this variation. This explicitly demonstrates the consistency of the BPHZ-renormalization scheme for the self-consistent approximations: The operations of variation with respect to G and renormalization are commutative, i.e., one can construct the renormalized self-energy in two equivalent ways: The first uses the *un-renormalized* Φ -functional and defines the renormalized self-energy by applying the BPHZ-renormalization theorem to its diagrams which are defined by opening any line of the un-renormalized Φ -functional. In this way we have defined the renormalized self-energy in the previous section. Subsequently we renormalized the Γ -functional by substituting the thereby defined renormalized functions for the divergent vacuum-sub-diagrams with 2 and 4 external legs. After subtracting the pure vacuum contribution this leads to the finite renormalized functional (47).

As we have seen now, the second way to define the renormalized self-energy is to renormalize the Γ -functional first. Then the variation with respect to the *renormalized* propagator leads to the *renormalized* equations of motion and thus directly to the *renormalized* self-energy which shows the consistency of the local vacuum counter terms including all combinatorial factors for both, the generating functional and the equation of motion.

We like to clarify that the equations derived in this section for the in-matter parts of the propagator are all valid also in the general non-equilibrium case of quantum field theory provided the density operator at time t_0 amends a Wick decomposition. This is valid for statistical operators at initial time of the form [31]

$$\mathbf{R} = \frac{1}{Z} \exp\left(-\sum_k \alpha_k \mathbf{A}_k\right) \text{ with } Z = \text{Tr} \exp\left(-\sum_k \alpha_k \mathbf{A}_k\right), \quad (52)$$

where the \mathbf{A}_k are one-particle operators. Our arguments for the renormalizability with α_k independent (i.e., state independent) local counter terms should hold, since the statistical operator is

normalized $\text{Tr } \mathbf{R} = 1$. Thus, any in-matter part of the propagator leads to similar reductions of the degree of divergence as for the Bose-Einstein distribution functions used here. This ensures that the power counting arguments for the non-vacuum parts are still valid. Note in particular that the real-time functional Γ has only functional meaning, namely as a tool to derive the equations of motion, since its value at the physical solution vanishes.

E. Renormalization of the thermodynamical potential

Using the thermodynamic part, i.e., the vertical branch, of the contour given in Fig. 1 the Γ -functional provides a finite value which indeed relates to the thermodynamic potential Ω . For the evaluation one uses the relationship between the Matsubara functions and the real-time functions given in Appendix A, Eq. (A21) for the thermodynamic trace Tr_{Th} which takes due account of the thermodynamic weights in the partition sum

$$\begin{aligned} \text{Tr}_{\text{Th}}\{h(p)\} &= \beta V \int \frac{d^d p}{(2\pi)^d} (h^{-+} + h^{+-}) \\ &= -2i\beta V \int \frac{d^d p}{(2\pi)^d} \text{sign}(p_0) \left(n(p_0) + \frac{1}{2} \right) \text{Im } h_R(p), \end{aligned} \quad (53)$$

where h^{-+} and h^{+-} are the Wightman functions and h_R is the retarded function of h . Furthermore

$$n(p_0) = \frac{1}{\exp(\beta|p_0|) - 1} \quad (54)$$

is the thermal Bose-Einstein factor resulting from the summation over Matsubara frequencies expressed in terms of complex contour integrals cf. (A21). For this thermal contour \mathcal{C}_{Th} closed diagrams as those of Φ and Γ also attain a finite value. In this case the rule is, first to omit one of the momentum integrations, which in this way defines a two-point function. Its renormalized retarded value can be calculated according to the above used real-time contour rules. Subsequently one applies (53) for the final integration. We show now that after renormalization of the two-point function we need only to subtract the overall vacuum divergence inherent in this final integral.

To obtain this result we have to go back to the regularized expression for the *un-renormalized* effective potential (8) which relates to the thermodynamical potential via

$$\Omega^{(\text{reg})}(T) = -T\Gamma_{\text{Th}}^{(\text{reg})} \quad (55)$$

where

$$\begin{aligned} \Gamma_{\text{Th}}^{(\text{reg})}(T) &= \Gamma_{\text{Th}}^{(\text{vac,reg})}(T) + \Gamma_{\text{Th}}^{(\text{matter,reg})}(T) \quad \text{with} \\ \Gamma_{\text{Th}}^{(\text{vac,reg})}(T) &= \text{Tr}_{\text{Th}} \left[\frac{i}{2} \ln \left(-\frac{M^2}{G^{\text{vac}}} \right) + \frac{i}{2} \Sigma^{\text{vac}} G^{\text{vac}} \right] + \Phi_{\text{Th}}^{\text{vac}}(T), \\ \Gamma_{\text{Th}}^{(\text{matter,reg})}(T) &= \frac{i}{2} \text{Tr}_{\text{Th}} \left[G^{(\text{r})} \Sigma^{(\text{matter})} - \frac{1}{2} G^{(\text{vac})} \Sigma^{(\text{r})} G^{(\text{vac})} \Sigma^{(\text{r})} - \sum_{k=3}^{\infty} \frac{(G^{(\text{vac})} \Sigma^{(\text{matter})})^k}{k} \right] \\ &\quad + \frac{1}{4} G^{(\text{r})} \Lambda^{(\text{vac})} G^{(\text{r})} + \Phi_{\text{Th}}^{(\text{r})}(T). \end{aligned} \quad (56)$$

Here the subscript Th specifies the quantities resulting from the thermal trace. The matter part results from form (47). Subsequently one replaces all quantities by their renormalized ones (denoted

by a bar across the functions) and cancels the overall divergence by subtracting the $T \rightarrow +0$ value

$$\begin{aligned}\Omega^{(\text{ren})}(T) &= -T \left(\bar{\Gamma}_{\text{Th}}^{(\text{reg})}(T) - \bar{\Gamma}_{\text{Th}}^{(\text{reg})}(+0) \right) \\ &= -T \left(\bar{\Gamma}_{\text{Th}}^{(\text{vac,reg})}(T) - \bar{\Gamma}_{\text{Th}}^{(\text{vac,reg})}(+0) + \bar{\Gamma}_{\text{Th}}^{(\text{matter,reg})}(T) \right).\end{aligned}\quad (57)$$

This procedure is legitimate as long as the new loops due to the final thermodynamical trace (53) do not induce new subdivergences, but only overall divergences. It is obvious that for the matter part the final trace loop involving the factor $n + 1/2$ is completely convergent, since all loops are regular once Λ is renormalized. For the two vacuum terms the component proportional to the factor $1/2$ in the thermal trace (53) cancel out, such that all terms are proportional to $n(p_0)$ which cuts off the p_0 integration, while the momentum integrations are also limited due to the vacuum thresholds: the imaginary parts of vacuum functions are zero for $p^2 < m^2$, where m is the mass of the stable vacuum particle. Thus also these final loop integrals are finite, defining a finite thermodynamical potential.

The vacuum part essentially determines the kinetic energy part of Ω as can be seen from the most simple example of an ideal gas. Here of course all self-energies and Φ are vanishing, the retarded propagator at finite temperature is $D_R(p) = [p^2 - m^2 + i\eta\sigma(p_0)]^{-1}$ and the renormalization is done by subtracting the pure vacuum part. Thus, the free thermodynamical potential becomes

$$\Omega^{(\text{id. gas})} = -V \int \frac{d^4l}{(2\pi)^4} n(l_0) \pi \Theta(l^2 - m^2), \quad (58)$$

which can be brought to a more familiar form by an integration by parts

$$\Omega^{(\text{id. gas})} = -pV = V \int \frac{d^3\vec{l}}{(2\pi)^3} \ln \left[1 - \exp \left(-\beta \sqrt{\vec{l}^2 + m^2} \right) \right]. \quad (59)$$

IV. CONCLUSIONS AND OUTLOOK

For the example of ϕ^4 -theory we have shown that self-consistent Dyson resummations based on a Φ -derivable scheme can be renormalized with local counter terms defined on the self-consistently determined vacuum level. This result was obtained with help of Weinberg's power counting theorem and using the BPHZ-renormalization scheme with the usual modifications for finite temperature diagram rules, which can be summarized in the simple rule that the "contraction boxes" defining the counter terms have to exclude sub-diagrams which contain any temperature line.

The hidden subdivergence structure of the self-consistent scheme has been resolved. This leads to a Bethe-Salpeter equation for the vacuum four-point function compatible with the chosen Φ -approximation, which we have renormalized. The method is free of pinch singularities. Closed equations could be formulated which resum the non-perturbative structure of both, the equations of motion, i.e., the self-energies and also the non-perturbative counter-term structure. The complexity of these equations is comparable to standard Dyson resummation schemes and therefore in principle does not imply new techniques. First numerical applications, which include the construction of the BS-kernel, the solution of the half-sided four-point function and thus the renormalized self-energies up to the self-consistent sunset self-energy, are presented in a second paper [26]. The renormalization of the generating functional Γ , c.f. sect. IIID, shows that the derivation and thus the renormalized in-matter equations of motion equally apply to the general non-equilibrium case.

This also proves that there is no arbitrariness in studying the in-medium modifications of model parameters like the mass and the coupling constants within this class of approximation schemes: It is sufficient to adjust them in the vacuum, for instance by fitting them to scattering

data, in order to predict without ambiguity how they change in the dense and hot medium: *The in-medium modifications are ruled completely by the model and its vacuum parameters alone, no further assumptions need to be made.*

Although demonstrated for the ϕ^4 -theory, the method is in principle general, since the derivation only relies on the analytic and asymptotic form of the propagators. In particular the renormalization of hidden overlapping divergences in the logarithmically divergent Bethe-Salpeter equations is general. Still, there is a number of restrictions of the self consistent Dyson resummation within the Φ -derivable scheme, which concerns global and local symmetries and the corresponding conservation laws and Ward-Takahashi identities.

The Φ -functional formalism only ensures the conservation laws for the expectation values of charges associated with the symmetry by Noether's theorem. However, in general the Ward-Takahashi identities are violated for the self-energy and higher vertex functions. Heuristically the problem can be traced back to the violation of crossing symmetry by the self-consistent scheme: Our derivation shows that the self-consistent solution of the self-energy involves Bethe-Salpeter ladder resummations of the four-point function, but only in the s -channel. The crossing symmetric t - and u - channel contributions to the four-point function are not included.

The symmetry properties of the $\Gamma[\varphi, G]$ -functional were already investigated by us with help of the here applied path-integral method [32, 33]. We show that it is always possible to define a *non-perturbative* approximation to the effective action $\Gamma_{\text{eff}}[\varphi]$ which respects linearly realized symmetries of the classical action provided the symmetry is not anomalously broken. The self-energy and higher vertex functions defined from this improved approximation action formalism then fulfill the Ward-Takahashi identities of the underlying symmetry. As a result the effective action Γ_{eff} enforces that additional t - and u -channel Bethe-Salpeter resummations are needed to restore the crossing symmetry together with the Ward-Takahashi identities for the self-energy and the vertex functions. However, these vertex functions are not self-consistently calculated and thus some problems remain also within this approximation: For instance in the case of the linear sigma-model the $O(N)$ -symmetry is restored for the vertex-functions and the Goldstone-modes become massless. Yet, the phase transition from the Nambu-Goldstone phase at low temperatures to the Wigner-Weyl phase at high temperatures results to be of 1st order rather than 2nd order [9].

In the case of a local gauge symmetry the problems become even more intricate: Self-consistent schemes beyond the classical field level for the gauge fields generally violate local gauge symmetries for the same reasons as for global symmetries. However, this immediately causes the excitation of spurious modes of the gauge fields which leads to violation of the unitarity of the S-matrix, the positive definiteness of the statistical operator and the causality structure of Green's functions. Nevertheless a gauge invariant effective action Γ_{eff} within a background-field approach can be formulated which provides gauge covariant polarization functions [17].

From a practical point of view the problem remains to calculate the self-consistent propagators needed for the symmetry-restoring Bethe-Salpeter resummation, which presently can only be solved in simple cases (RPA bubble resummation). In [17] we have presented a workaround in terms of a suitably chosen projection method onto the physical (transverse) degrees of freedom of the gauge-field polarization tensor. This procedure, of course, does not lead to a full restoration of local gauge theory but to causal Green's functions and current conservation within the self-energies of matter-fields. Alternative methods are restricted to the approximate solution of the self-consistent equations of motion, e.g., in the sense of a Hard thermal loop approximation [19] or to a systematic expansion in terms of the coupling constant or \hbar [18].

The proof of the renormalizability of Φ -derivable approximations opens a broad range of perspectives for effective field theory model applications describing the non-perturbative in-medium properties of particles in dense or finite-temperature matter with model parameters fixed at the vacuum level. Further applications point towards the appropriate renormalization of non-equilibrium

transport equations [16], where in particular the drift terms, which determine the equation of state, involve the real part of the self-energies which generally need renormalization.

Acknowledgments

We are grateful to G. E. Brown, P. Danielewicz, B. Friman, Yu. Ivanov, E. E. Kolomeitsev, M. Lutz, M. A. Nowak and D. Voskresensky for fruitful discussions and suggestions at various stages of this work.

APPENDIX A: ANALYTICAL PROPERTIES OF GREEN'S FUNCTIONS

In this appendix we summarize briefly the analytic properties of Green's functions of neutral bosons needed in the main part of the paper. This is most easily done by switching to the operator formalism in the Heisenberg picture. By definition we have for a hermitian scalar field operator

$$iG(x) = \frac{1}{Z} \text{Tr} \exp(-\beta \mathbf{H}) \phi(x) \phi(0) := \langle \mathcal{T}_{\mathcal{C}} \phi(x) \phi(0) \rangle_{\beta}. \quad (\text{A1})$$

For x^0 on the vertical part, i.e., $x^0 = -i\tau$ with $0 \leq \tau \leq \beta$ we obtain the Matsubara Green's function

$$G_M(\tau, \vec{x}) = G^{+-}(-i\tau, \vec{x}), \quad (\text{A2})$$

where one has to understand the analytic continuation of the real-time Wightman function G^{+-} on the right hand side. It is important to keep in mind that (A2) is only valid when the first time argument x^0 in (A1) is on the vertical part of the contour while the second one is at $x^0 = 0$. If both arguments of the fields are located on the vertical part according to (A1) one has to use the time ordering along the imaginary time axis.

Since the order of the operators under the trace in (A1) can be changed cyclically the real-time Wightman functions G^{+-} and G^{-+} are related through

$$G^{+-}(x^0 - i\beta, \vec{x}) = G^{-+}(x^0, \vec{x}), \quad (\text{A3})$$

where x^0 is a real-time argument on the contour and on the right hand side one has to understand the analytic continuation. One should keep in mind that only the Wightman functions are analytically continueable, not the time ordered or anti-time ordered of the upper or lower real-time branch, since they contain step functions from the time ordering operator $\mathcal{T}_{\mathcal{C}}$. The same holds true for the imaginary-time ordered Matsubara Green's function.

As in vacuum quantum field theory due to translation invariance it is customary to use the energy-momentum representation of Green's functions. For the real-time propagators we have the usual description

$$G^{ij}(x) = \int \frac{d^d p}{(2\pi)^d} \exp(-ipx) G^{ij}(p). \quad (\text{A4})$$

We write down the formalism for arbitrary space-time dimensions since all considerations do not depend on it and we need it to obtain well defined non-renormalized quantities in the sense of dimensional regularization. The periodic boundary condition (A3) translates into the *Kubo-Martin-Schwinger-condition* (KMS) for the Fourier transformed Green's functions:

$$G^{-+}(p) = \exp(-\beta p_0) G^{+-}(p). \quad (\text{A5})$$

The Matsubara Green's function is only defined for imaginary times $-i\tau$ with $0 \leq \tau \leq \beta$. Thus the momentum representation with respect to the time component is a Fourier series with period β according to (A3) rather than a Fourier integral:

$$G_M(x) = \frac{1}{\beta} \sum_{n=-\infty}^{\infty} \int \frac{d^{d-1}\vec{p}}{(2\pi)^{d-1}} \exp(-ipx) G_M(ip_0, \vec{p})|_{p^0=\omega_n} \text{ with } \omega_n = \frac{2\pi}{\beta}n. \quad (\text{A6})$$

Herein $x^0 = -i\tau$ with $0 \leq \tau \leq \beta$. Using the inverse Fourier transform, Eq. (A4), and the KMS-condition (A5) we find the *spectral representation*

$$G_M(i\omega_n, \vec{p}) = i \int \frac{dp_0}{2\pi} \frac{\rho(p_0, \vec{p})}{p_0 - i\omega_n} \text{ with } \rho(p) = i[G^{+-}(p) - G^{-+}(p)]. \quad (\text{A7})$$

This shows that the Matsubara propagator is completely represented by the real-time function ρ . With help of this we define the analytically continued propagator by

$$G_c(k) = - \int \frac{dp_0}{2\pi} \frac{\rho(p_0, \vec{k})}{p_0 - k_0}. \quad (\text{A8})$$

It can contain singularities only on the real axis. Using the Fourier transformation (A4) for the limits to the real axis from above and below we obtain:

$$G_c(p_0 \pm i\eta, \vec{p}) = G_{R/A}(p), \quad p_0 \in \mathbb{R} \quad (\text{A9})$$

with the *retarded and advanced* Green's functions

$$G_{R/A}(x) = \mp i\Theta(\pm t) \langle [\phi(x), \phi(0)]_{\mp} \rangle_{\beta}. \quad (\text{A10})$$

From this we find immediately

$$G_R(x) = G_A^*(-x) \Rightarrow G_R(p) = G_A^*(p), \quad \rho(p) = -2 \text{Im} G_R(p) = -\rho(-p) \quad (\text{A11})$$

and from (A8) and the analyticity of $G_c(p)$ in the upper complex p_0 -plane it follows that

$$\sigma(p_0)\rho(p) \geq 0. \quad (\text{A12})$$

For later use we note the momentum space properties

$$G^{--} + G^{++} = G^{+-} + G^{-+}, \quad G_R = G^{--} - G^{-+}, \quad G_A = G^{--} - G^{+-}, \quad G_M = -iG_c(i\omega_n) \quad (\text{A13})$$

which follow immediately from (A1), (A7), and (A10).

We also make use of the expressions for the real-time Green's functions in terms of the retarded Green's function, which follow immediately from (A7,A11,A18):

$$iG^{--}(p) = iG_R(p) + [\Theta(-p_0) + n(p_0)]\rho(|p_0|, \vec{p}), \quad (\text{A14})$$

$$iG^{++}(p) = [\Theta(p_0) + n(p_0)]\rho(|p_0|, \vec{p}) - iG_R(p), \quad (\text{A15})$$

$$iG^{-+}(p) = [\Theta(-p_0) + n(p_0)]\rho(|p_0|, \vec{p}), \quad (\text{A16})$$

$$iG^{+-}(p) = [\Theta(p_0) + n(p_0)]\rho(|p_0|, \vec{p}). \quad (\text{A17})$$

and the *Bose-Einstein distribution defined as*

$$n(p_0) = \frac{1}{\exp(\beta|p_0|) - 1}. \quad (\text{A18})$$

All relations given above for the Green's functions G directly apply to any two-point function given by local field operators $ih(x, y) = \langle \mathcal{T}_{\mathcal{C}_{\mathbb{R}}} \mathbf{H}(x) \mathbf{H}(y) \rangle$, e.g., the self-energy.

Real-time contour integrations and traces of translationally invariant two-point functions

$$\begin{aligned} C(x, y) &= \int_{\mathcal{C}_{\mathbb{R}}} dz A(x, z) B(z, y), \quad x, y, z \in \mathcal{C}_{\mathbb{R}} \\ \text{Tr}_{\mathbb{R}} C &= \int_{\mathcal{C}_{\mathbb{R}}} dx C(x, x) \end{aligned} \tag{A19}$$

transcribe to

$$\begin{aligned} C(p)^{ij} &= \sum_{kl} A(p)^{ik} \sigma_{kl} B(p)^{lj}, \quad \sigma = \text{diag}(1, -1), \quad i, j, k, l \in \{-, +\} \\ \text{Tr}_{\mathbb{R}} C &= \sum_{ij} \int \frac{d^d p}{(2\pi)^d} C(p)^{ij} \sigma_{ij} \end{aligned} \tag{A20}$$

in contour momentum-space representation.

We close this appendix by citing the formula for summation over the Matsubara frequencies needed when calculating quantities related to the vertical branch of the contour [24]. In this paper we use this to calculate the thermodynamical potential

$$\begin{aligned} \text{Tr}_{\text{Th}} h(p) &:= \beta V \frac{1}{i\beta} \sum_{n=-\infty}^{\infty} \int \frac{d^{d-1} \vec{p}}{(2\pi)^{d-1}} h(i\omega_n, \vec{p}) \\ &= -\beta V \int \frac{d^d p}{(2\pi)^d} \left[\frac{1}{2} + n(p_0) \right] \{ h[p_0 + i\eta\sigma(p_0)] - h[p_0 - i\eta\sigma(p_0)] \} \\ &= \beta V \int \frac{d^d p}{(2\pi)^d} (h^{-+} + h^{+-}) \end{aligned} \tag{A21}$$

In (A21) we have assumed that the function h is analytic below and above the real axis and that the trace exists. Usually this is only the case for the regularized or the renormalized functional traces. It is also clear that due to the exponential damping from the Bose-Einstein distribution (A18) this part of the integral has a superficial degree of divergence reduced by 1 compared to the first part which is not damped by an n -factor.

Eq. (A21) shows that the thermodynamical potential can be calculated from real-time quantities since the analytic continuation of the Matsubara Green's function needed on the right hand side is unique and can be obtained from the retarded Green's function as well c.f. (A9). For a more detailed analysis of the analytic properties see also [25] and for the general case of Wigner function representations in the non-equilibrium context [11].

-
- [1] J. Luttinger and J. Ward, Phys. Rev. **118**, 1417 (1960), URL http://prola.aps.org/abstract/PR/v118/i5/p1417_1.
 - [2] T. D. Lee and C. N. Yang, Phys. Rev. **117**, 22 (1961), URL http://prola.aps.org/abstract/PR/v117/i1/p22_1.
 - [3] G. Baym and L. Kadanoff, Phys. Rev. **124**, 287 (1961), URL http://prola.aps.org/abstract/PR/v124/i2/p287_1.
 - [4] G. Baym, Phys. Rev. **127**, 1391 (1962), URL http://prola.aps.org/abstract/PR/v127/i4/p1391_1.

- [5] M. Cornwall, R. Jackiw, and E. Tomboulis, Phys. Rev. **D10**, 2428 (1974), URL http://prola.aps.org/abstract/PRD/v10/i8/p2428_1.
- [6] J. Schwinger, J. Math. Phys **2**, 407 (1961).
- [7] L. Keldysh, ZhETF **47**, 1515 (1964), [Sov. Phys JETP **20** 1965 1018].
- [8] S. A. Chin, Ann. Phys. **108**, 301 (1977).
- [9] G. Baym and G. Grinstein, Phys. Rev. **D15**, 2897 (1977), URL http://prola.aps.org/abstract/PRD/v15/i10/p2897_1.
- [10] A. F. Bielajew and B. D. Serot, Annals of Physics **156**, 215 (1984).
- [11] Y. B. Ivanov, J. Knoll, and D. N. Voskresensky, Nucl. Phys. **A657**, 413 (1999), URL <http://arXiv.org/abs/hep-ph/9807351>.
- [12] J. Knoll, Prog. Part. Nucl. Phys. **42**, 177 (1999), URL <http://arXiv.org/abs/nucl-th/9811099>.
- [13] Y. B. Ivanov, J. Knoll, and D. N. Voskresensky, Nucl. Phys. **A672**, 313 (2000), URL <http://arXiv.org/abs/nucl-th/9905028>.
- [14] S. Leupold, Nucl. Phys. **A672**, 475 (2000).
- [15] L. Kadanoff and G. Baym, *Quantum Statistical Mechanics*, A Lecture Note and Preprint Series (The Benjamin/Cummings Publishing Company, 1961).
- [16] Y. I. Knoll, J. and D. Voskresensky, Annals of Physics (2001), in press, URL <http://arXiv.org/abs/nucl-th/0102044>.
- [17] H. van Hees and J. Knoll, Nucl. Phys. **A683**, 369 (2001), URL <http://arXiv.org/abs/hep-ph/0007070>.
- [18] J. P. Blaizot, E. Iancu, and A. Rebhan, Phys. Rev. **D63**, 065003 (2001), URL <http://arXiv.org/abs/hep-ph/0005003>.
- [19] A. Peshier, Phys. Rev. **D63**, 105004 (2001), URL <http://arXiv.org/abs/hep-ph/0011250>.
- [20] S. Weinberg, Phys. Rev. **118**, 838 (1960), URL http://prola.aps.org/abstract/PR/v118/i3/p838_1.
- [21] N. N. Bogoliubov and O. S. Parasiuk, Acta Math. **97**, 227 (1957).
- [22] W. Zimmermann, Commun. Math. Phys. **15**, 208 (1969).
- [23] J. C. Collins, *Renormalization* (Cambridge University Press, Cambridge, New York, Melbourne, 1986).
- [24] J. I. Kapusta, *Finite-Temperature Field Theory* (Cambridge University Press, Cambridge, New York, Melbourne, 1989).
- [25] M. LeBellac, *Thermal Field Theory* (Cambridge University Press, Cambridge, New York, Melbourne, 1996).
- [26] H. van Hees and J. Knoll, Phys. Rev. D (2001), to be submitted.
- [27] N. P. Landsmann and C. G. van Weert, Physics Reports **145**, 141 (1987).
- [28] F. Gelis, Z. Phys. **C70**, 321 (1996), URL <http://arXiv.org/abs/hep-ph/9412347>.
- [29] C. Itzykson and J.-B. Zuber, *Quantum Field Theory* (McGraw-Hill Book Company, New York, 1980).
- [30] W. Zimmermann, Lectures on Elementary Particles and Quantum Field Theory. M.I.T. Press p. 397 (1970).
- [31] P. Danielewicz, Annals of Physics **152**, 239 (1984).
- [32] H. van Hees, Ph.D. thesis, TU Darmstadt (2000), URL <http://elib.tu-darmstadt.de/diss/000082/>.
- [33] H. van Hees and J. Knoll, Phys. Rev. D (2001), in preparation.